



Calhoun: The NPS Institutional Archive
DSpace Repository

Reports and Technical Reports

All Technical Reports Collection

1999-01

Convolution methods for mathematical problems in biometrics

Frenzen, Christopher L.

Monterey, California. Naval Postgraduate School

<http://hdl.handle.net/10945/24439>

Downloaded from NPS Archive: Calhoun



Calhoun is the Naval Postgraduate School's public access digital repository for research materials and institutional publications created by the NPS community. Calhoun is named for Professor of Mathematics Guy K. Calhoun, NPS's first appointed -- and published -- scholarly author.

Dudley Knox Library / Naval Postgraduate School
411 Dyer Road / 1 University Circle
Monterey, California USA 93943

<http://www.nps.edu/library>

NPS-MA-99-001

NAVAL POSTGRADUATE SCHOOL Monterey, California



CONVOLUTION METHODS FOR MATHEMATICAL PROBLEMS IN BIOMETRICS

by

Christopher Frenzen

January 15, 1999

Approved for public release; distribution is unlimited.

Prepared for: Biometrics Identification
College of Engineering
San Jose State University
One Washington Square
San Jose, CA 95192-0205

1 9 9 9 0 2 1 8 0 8 4

REPORT DOCUMENTATION PAGEForm approved
OMB No 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE January 15, 1999	3. REPORT TYPE AND DATES COVERED March 1997 – September 1997	
4. TITLE AND SUBTITLE Convolution Methods for Mathematical Problems in Biometrics			5. FUNDING MIPR # NAFRLP62800493	
6. AUTHOR(S) Christopher L. Frenzen				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Postgraduate School Monterey, CA 93943			8. PERFORMING ORGANIZATION REPORT NUMBER NPS-MA-99-001	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Biometrics Identification College of Engineering San Jose State University One Washington Square San Jose, CA 95192-0205			10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES Approved for public release; distribution is unlimited.				
12a. DISTRIBUTION/AVAILABILITY STATEMENT			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words.) Estimation of the impostor probability density function from the inter-template and template-sample probability density functions involves a multi-dimensional convolution of the latter density functions. We assume isotropy of the probability distribution functions and use the Fourier transform to express the convolution as a one-dimensional integral with a kernel given by a spherical Bessel function.				
14. SUBJECT TERMS Biometrics, Convolution			15. NUMBER OF PAGES 6	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED	18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	20. LIMITATION OF ABSTRACT	

NAVAL POSTGRADUATE SCHOOL
Monterey, California 93943-5000

RADM Robert C. Chaplin
Superintendent

R. Elster
Provost


This report was prepared for Biometrics Identification, College of Engineering, San Jose State University.

This report was prepared by:




Christopher L. Frenzen
Professor of Mathematics

Reviewed by:



Guillermo Owen, Chair
Department of Mathematics

Released by:



D. W. Netzer
Associate Provost and
Dean of Research

Convolution Methods for Mathematical Problems in Biometrics

C.L. Frenzen
Department of Mathematics
Naval Postgraduate School, Monterey, CA

1. Introduction. The problem we shall investigate can be formulated in the following way, due to Peter Bickel [1]: For $M + 1$ individuals K measurements $X_{ij}, i = 1, \dots, M + 1, j = 1, \dots, K$ are made. The X_{ij} are assumed to be vectors in \mathbf{R}^N .

We assume, following Bickel [1], that

$$X_{ij} = \mu_i + \varepsilon_{ij},$$

where μ_1, \dots, μ_{M+1} are individual effects vectors from \mathbf{R}^N which are i.i.d. (independently and identically distributed) with a distribution which is spherically symmetric about some μ in \mathbf{R}^N . The ε_{ij} are i.i.d. independently of the μ_i and their distribution is spherically symmetric about 0.

The $K(M + 1)X_{ij}$ s represent K biometric measurements made on $M + 1$ individuals; however the measuring device actually records only the data $|\bar{X}_i - \bar{X}_{i'}|$ and $|X_{M+1,j} - \bar{X}_{M+1}|$, $1 \leq i < i' \leq M, 1 \leq j \leq k$, where $|x|$ is the length of x and

$$\bar{X}_i = \frac{1}{K} \sum_{j=1}^K X_{ij}$$

is the centroid of the K measurements taken on the the i th individual.

Let p be the density of $|\bar{X}_1 - \bar{X}_2|$, q the density of $|X_{M+1,1} - \bar{X}_{M+1}|$, and r the density of $|X_{M+1,1} - \bar{X}_1|$. The basic problem is to estimate r given estimates \hat{p}, \hat{q} for p and q . As p and q are densities on $[0, \infty)$, the problem of estimating \hat{p} and \hat{q} is a standard problem in estimation theory which we do not consider further.

2. Bickel's Approach. What is the relationship between p , q , and r ? Following Bickel [1], we write

$$X_{M+1,1} - \bar{X}_1 = \varepsilon_{M+1,1} + \mu_{M+1} - \mu_1 - \bar{\varepsilon}_1, \quad (1)$$

$$\bar{X}_{M+1} - \bar{X}_1 = \bar{\varepsilon}_{M+1} + \mu_{M+1} - \mu_1 - \bar{\varepsilon}_1, \quad (2)$$

$$X_{M+1,1} - \bar{X}_{M+1} = \varepsilon_{M+1,1} - \bar{\varepsilon}_{M+1}. \quad (3)$$

Note that addition of equations (2) and (3) yields equation (1). Further, the quantities on the left sides of (2) and (3) have spherically symmetric distributions. As the lengths of the left sides of (2) and (3) have densities p and q respectively, and the length of the left side of (1) is r , it seems that what is required is a formula for the density of the length of the convolution of two spherically symmetric (about 0) distributions given the densities of their lengths. However, as Bickel pointed out in [1], this is not quite correct since the use of the convolution assumes the independence of the densities p and q , and these two densities are not generally independent, since in the right sides of (2) and (3) the terms $\varepsilon_{M+1,1} - \bar{\varepsilon}_{M+1}$ and $\bar{\varepsilon}_{M+1}$ are only uncorrelated, and not in general independent unless ε_{11} has a Gaussian distribution.

However, for large K , we follow Bickel's argument in [1] to show that the terms $\varepsilon_{M+1,1} - \bar{\varepsilon}_{M+1}$ and $\bar{\varepsilon}_{M+1}$ are independent. Let

$$Z_K = (K-1)^{-\frac{1}{2}} \sum_{j=2}^K \varepsilon_{M+1,j}, \quad (4)$$

and

$$U_K = \varepsilon_{M+1,1} \left(1 - \frac{1}{K}\right). \quad (5)$$

Then, Z_K and U_K are independent and, if K is large,

$$\varepsilon_{M+1,1} - \bar{\varepsilon}_{M+1} = U_K - \frac{(K-1)^{\frac{1}{2}}}{K} Z_K \approx U_K \quad (6)$$

and

$$K^{\frac{1}{2}} \bar{\varepsilon}_{M+1} = \left(\frac{K-1}{K}\right)^{\frac{1}{2}} Z_K + U_K \frac{K^{\frac{1}{2}}}{(K-1)} \approx Z_K. \quad (7)$$

Thus, for large K , to a first approximation it is possible to ignore the dependence between (6) and (7). The terms U_K and Z_K on the right sides of (6) and (7) respectively are the first terms in an asymptotic expansion for large K of the left sides of those equations.

3. Convolution and Fourier Transform. Let U, V be random vectors in \mathbf{R}^N which are independent and have isotropic distributions with densities

f_1, f_2 respectively. We let g_1, g_2 be the corresponding densities on \mathbf{R}^+ of the lengths of U and $V, |U|, |V|$. Further, let f be the density of $U + V$ and g be the density of $|U + V|$. Then by the independence of $U, V,$

$$f(y) = \int_{\mathbf{R}^N} f_2(y - x) f_1(x) dx. \quad (8)$$

Our interest is to determine a formula for g in terms of g_1 and g_2 . To this end, we introduce the Fourier transform. If f is absolutely integrable on \mathbf{R}^N , then the Fourier transform of f is defined by

$$\hat{f}(t) = \frac{1}{(2\pi)^{N/2}} \int_{\mathbf{R}^N} f(y) e^{it \cdot y} dy, \quad (9)$$

where the N vector $t = (t_1, t_2, \dots, t_N)$ and $t \cdot y = t_1 y_1 + \dots + t_N y_N$. If $f(y)$ is spherically symmetric, i.e., $f(y)$ is a function of $r = |y|$ only, say $f(y) = h(r)$, then its Fourier transform $\hat{f}(t)$ is also spherically symmetric; more specifically, we have

$$\hat{f}(t) = \rho^{(2-N)/2} \int_0^\infty r^{N/2} h(r) J_{(N-2)/2}(\rho r) dr, \quad (10)$$

where $\rho = |t|$ and $J_\nu(r)$ denotes the Bessel function of the first kind of order ν . For a proof of this formula, we refer to Schwartz[2]. (Note that we have introduced the same letter r for $|y|$ as we used for the density of the length of the left side of (1). From the context, there should be no confusion as to which meaning for r is intended.) The Fourier transform of the convolution in (8) yields

$$\hat{f}(t) = \hat{f}_2(t) \hat{f}_1(t), \quad (11)$$

and combining (10) and (11) gives

$$\hat{f}(t) = \rho^{2-N} \int_0^\infty r^{N/2} h_2(r) J_{(N-2)/2}(\rho r) dr \int_0^\infty r^{N/2} h_1(r) J_{(N-2)/2}(\rho r) dr, \quad (12)$$

where the functions h_1, h_2 in (12) are defined by

$$h_1(r) = f_1(y), \quad h_2(r) = f_2(y), \quad (13)$$

with $r = |y|$ since the distributions defined by f_1, f_2 are isotropic.

Now if U is an isotropically distributed N vector with density f on \mathbf{R}^N , and g is the corresponding density on \mathbf{R}^+ of the length $|U|$, where $|x|^2 =$

$\sum_{i=1}^N x_i^2, x = (x_1, \dots, x_N)$, then the relationship between the densities f and g is given by

$$f(y) = |y|^{-(N-1)} c_N^{-1} g(|y|), \quad (14)$$

where c_N , the surface 'area' of the unit sphere in \mathbf{R}^N , is

$$c_N = \frac{2\pi^{N/2}}{\Gamma(\frac{N}{2})}. \quad (15)$$

Hence, upon substituting

$$h_1(r) = f_1(y) = r^{-(N-1)} c_N^{-1} g_1(r), \quad (16)$$

$$h_2(r) = f_2(y) = r^{-(N-1)} c_N^{-1} g_2(r),$$

into (12), we have

$$\hat{f}(t) = \rho^{2-N} c_N^{-2} \int_0^\infty r^{1-N/2} g_2(r) J_{(N-2)/2}(\rho r) dr \int_0^\infty r^{1-N/2} g_1(r) J_{(N-2)/2}(\rho r) dr. \quad (17)$$

If the dimension N of the space the measurement vectors X_{ij} belong to is even, say $N = 2m$, then $J_{(N-2)/2}(\rho r) = J_{m-1}(\rho r)$ is a Bessel function of the first kind of integer order. If N is odd, say $N = 2m + 1$, then $J_{(N-2)/2}(\rho r) = J_{m-1/2}(\rho r)$ is a Bessel function of the first kind of fractional order, and is closely related to the Spherical Bessel function of the first kind $j_n(z)$, defined by

$$j_n(z) = \sqrt{\frac{\pi}{2z}} J_{n+1/2}(z). \quad (18)$$

More detailed information about Bessel functions may be in found in Abramowitz and Stegun [3].

4. Convergence of the Integrals. We now discuss convergence of the integrals in (17). Both integrals are functions of the variable $\rho = |t|$, hence $\hat{f}(t)$ is a spherically symmetric function of the transform variable t . This means that the inverse Fourier transform of $\hat{f}(t)$, i.e. $f(y)$, can also be obtained as a one-dimensional integral with a Bessel function kernel.

For fixed ρ and small r , $J_{(N-2)/2}(\rho r) = O(r^{(N-2)/2})$. Since we expect $g_i^{(j)}(0) = 0$ for $i = 1, 2$ for $0 \leq j \leq N - 1$ (see Bickel [1]), it follows that both

integrals in (17) are convergent at the lower limit 0. At the upper limit ∞ , $J_{(N-2)/2}(\rho r) = O(r^{-1/2})$, and this by itself will not be enough to make the integral convergent. However the term $r^{1-N/2}$ will also make the integrals converge at the upper limit if N is sufficiently large. In practice the densities g_1, g_2 also tend to zero sufficiently rapidly to make the integrals converge at the upper limit. If these densities have compact support (i.e., are zero outside of a closed bounded subset of \mathbf{R}^+), then the integrals in (17) no longer have infinite upper limits. The integrals in (17) can be evaluated accurately and efficiently by standard numerical quadrature methods.

5. Inversion. Note that the right side of (17) is a function of $\rho = |t|$ only. Hence $\hat{f}(t)$ is spherically symmetric. Let $\hat{f}(t) = G(\rho)$. The inverse Fourier transform of $\hat{f}(t)$, i.e., $f(y)$ from (9), is defined by

$$f(y) = \frac{1}{(2\pi)^{N/2}} \int_{\mathbf{R}^N} \hat{f}(t) e^{-it \cdot y} dt. \quad (19)$$

It follows by analogy with (10) that

$$f(y) = r^{(2-N)/2} \int_0^\infty \rho^{N/2} G(\rho) J_{(N-2)/2}(\rho r) d\rho, \quad (20)$$

where $r = |y|$. The relationship (14) between the density f of an isotropically distributed N vector and the corresponding density g of its length then implies

$$g(r) = r^{N/2} c_N \int_0^\infty \rho^{N/2} G(\rho) J_{(N-2)/2}(\rho r) d\rho \quad (21)$$

where $G(\rho) = \hat{f}(t)$ is given by (17). Numerical evaluation of the integral in (21) proceeds similarly to the integrals in (17). With g_1, g_2 taken as p, q introduced at the end of section 1, and g taken as r (the density function for $|X_{M+1,1} - \bar{X}_1|$, not $|y|$), (17) and (21) together give the density r in terms of p and q .

6. Conclusions. We have shown that for large K , the left sides of (2) and (3) are approximately independent. Under this approximation, we

derived an expression for the density r of the length of the left side of (1) in terms of the densities p, q of the lengths of the left sides of (2) and (3). This result is contained in equations (17) and (21) of the previous section.

Future work could include determining corrections to the above result for a finite number of measurements K , and practical numerical implementation of the above result.

References

- 1.) NSA SAG Problem 97-2-1 Solution, Peter Bickel.
- 2.) *Mathematics for Physical Sciences*, L. Schwartz, Addison-Wesley, Reading, MA, 1966, pp. 201-203.
- 3.) *Handbook of Mathematical Functions*, A. Abramowitz and I. Stegun, National Bureau of Standards, 1964.

DISTRIBUTION LIST

Director Defense Tech Information Center Cameron Station Alexandria, VA 22314	(2)
Research Office Naval Postgraduate School Monterey, CA 93943	(1)
Library Naval Postgraduate School Monterey, CA 93943	(2)
Professor C. L. Frenzen Department of Mathematics Naval Postgraduate School Monterey, CA 93943	(5)
Dr. James L. Wayman Research Director, Biometrics Identification College of Engineering San Jose State University One Washington Square San Jose, CA 95192-0205	(2)