



**Calhoun: The NPS Institutional Archive**  
**DSpace Repository**

---

Reports and Technical Reports

All Technical Reports Collection

---

1974-07

# Minimal point cubatures of precision seven for symmetric planar regions: Revisited

Franke, Richard H.

Monterey, California. Naval Postgraduate School

---

<http://hdl.handle.net/10945/29973>

*Downloaded from NPS Archive: Calhoun*



Calhoun is a project of the Dudley Knox Library at NPS, furthering the precepts and goals of open government and government transparency. All information contained herein has been approved for release by the NPS Public Affairs Officer.

**Dudley Knox Library / Naval Postgraduate School**  
**411 Dyer Road / 1 University Circle**  
**Monterey, California USA 93943**

<http://www.nps.edu/library>

NPS-53Fe74071

# NAVAL POSTGRADUATE SCHOOL

## Monterey, California



MINIMAL POINT CUBATURES OF PRECISION SEVEN FOR  
SYMMETRIC PLANAR REGIONS, REVISITED

Richard Franke

July 17, 1974

Technical Report For Period  
October 1973-December 1973

Approved for public release; distribution unlimited

Prepared for:  
Chief of Naval Research, Arlington, Virginia 22217

FEDDOCS  
D 208.14/2:NPS-53Fe74071

NAVAL POSTGRADUATE SCHOOL  
Monterey, California

Rear Admiral Isham Linder  
Superintendent

Jack R. Borsting  
Provost

The work reported herein was supported in part by the Foundation Research Program of the Naval Postgraduate School with funds provided by the Chief of Naval Research.

Reproduction of all or part of this report is authorized.

This report was report was prepared by:

L. D. KOVACH  
(Acting) Chairman of Mathematics

JOHN M. WOZENCRAFT ✓  
Dean of Research ✓

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER NPS-53Fe74071A	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) MINIMAL POINT CUBATURES OF PRECISION SEVEN FOR SYMMETRIC PLANAR REGIONS, REVISITED.	5. TYPE OF REPORT & PERIOD COVERED Technical Report October 73-December 73	
	6. PERFORMING ORG. REPORT NUMBER	
7. AUTHOR(s) Richard Franke	8. CONTRACT OR GRANT NUMBER(s)	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Foundation Research Program Naval Postgraduate School Monterey California 93940	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 61152N; RR000-01; RR000-01-10; PO4-0001	
11. CONTROLLING OFFICE NAME AND ADDRESS Chief of Naval Research Arlington, Virginia 22217	12. REPORT DATE 17 July 1974	
	13. NUMBER OF PAGES 10	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	15. SECURITY CLASS. (of this report)  UNCLASSIFIED	
	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release ; distribution unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Minimal Point cubature, Cubature, Planar region, Polynomial precision, Orthogonal polynomials, Symmetric region		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)  The key theorem concerning a lower bound for the number of points required by cubature formulas of precision seven for certain symmetric planar regions is extended to arbitrary symmetric planar regions, with a simplified proof.		



## 1.0 Introduction

The purpose of this paper is to give a simplified proof of a theorem that was previously stated in a more restricted form [1]. In that paper it was shown how to construct 12 point cubature formulas having precision seven for symmetric planar regions. For fully symmetric regions and rectangular regions the construction cannot fail, and the resulting formulas use the minimum number of points. In the next section we establish 12 points as a lower bound for all symmetric (in each variable) planar regions. The existence of 12 point formulas has not been established for arbitrary symmetric regions. The notation of this paper is the same as that of [1].

## 2.0 Minimal Point Formulas

Let  $R$  be a symmetric (about each axis) planar region and let  $w(x,y)$  be symmetric in each variable. We consider cubature formulas of the form

$$(1) \quad \int_R wf \approx \sum_{k=1}^N A_k f(v_k)$$

As a consequence of the characterization theorem given by Stroud [2], or [3, p.111], we have the following result, also given in [1].

Proposition 1: Let there be given a formula of type (1), with precision 7, for a region in the plane. Suppose that  $N < 15$ . Then the points  $v_k$ ,  $k=1, \dots, N$ , are zeros of  $15 - N$  linearly independent polynomials of degree 4, each of which is orthogonal, over  $R$  with respect to  $w$ , to all polynomials of degree  $\leq 3$ .

Our approach, as before, is to show that for the regions under consideration, there do not exist five orthogonal polynomials of degree four with 10 common zeros, nor four with 11 common zeros.

Theorem 2: Let the cubature formula (1) have precision seven, where  $R$  and  $w$  are symmetric. Then  $N \geq 12$ . Proof: We first establish that no 10 point formula exists, that being the minimum possible for any region. Assume the contrary. Then the basic orthogonal polynomials, which have the form

$$\begin{aligned}
 P^{(4,0)} &= x^4 + a_4 x^2 + b_4 y^2 + c_4 \\
 P^{(3,1)} &= x^3 y + a_3 xy \\
 (2) \quad P^{(2,2)} &= x^2 y^2 + a_2 x^2 + b_2 y^2 + c_2 \\
 P^{(1,3)} &= xy^3 + a_1 xy \\
 P^{(0,4)} &= y^4 + a_0 x^2 + b_0 y^2 + c_0
 \end{aligned}$$

must have the ten nodes as common zeros. Inspection of  $P^{(3,1)}$  and  $P^{(1,3)}$  reveal that their common zeros are  $(\pm\alpha, \pm\beta)$ , where  $\alpha^2 = -a_3$  and  $\beta^2 = -a_1$ , and the axes. However,  $P^{(2,2)}$  has at most two finite zeros on each axis, hence  $P^{(3,1)}$ ,  $P^{(2,2)}$ , and  $P^{(1,3)}$  can have at most eight common zeros. Hence no ten point formula can exist.

We now show that no 11 point formula exists. If one did there would exist four linearly independent orthogonal polynomials of which the nodes would be common zeros. This would mean that all the  $v_k$  lie on some nontrivial linear combination of any two of the orthogonal polynomials (2). For some  $\mu_1$  and  $\lambda_1$  they lie on  $P_1 = \mu_1 P^{(3,1)} + \lambda_1 P^{(1,3)} = xy(\mu_1 x^2 + \lambda_1 y^2 + \mu_1 a_3 + \lambda_1 a_1)$ . Thus the nodes lie on the axes or the conic  $Q_2 = \mu_1 x^2 + \lambda_1 y^2 + \mu_1 a_3 + \lambda_1 a_1$ . We now consider two possibilities: (i) one of  $\mu_1$  or  $\lambda_1$  is zero, i.e., the nodes lie on  $P^{(3,1)}$  or  $P^{(1,3)}$ ; (ii)  $\mu_1 \lambda_1 \neq 0$ .

Case (i): We show this cannot happen. We assume the nodes lie on  $P^{(3,1)}$ , and the argument is dual if they lie on  $P^{(1,3)}$ . The nodes also lie on  $P_2 = \mu_2 P^{(4,0)} + \lambda_2 P^{(2,2)}$  for some  $\mu_2$  and  $\lambda_2$ , not both zero, and on



$P_3 = \mu_3 P^{(0,4)} + \lambda_3 P^{(2,2)}$  for some  $\mu_3$  and  $\lambda_3$ , not both zero. We see that

$P_2$  has at most two zeros on the  $x$  axis. The remaining nodes must lie on  $x^2 + a_3$  and  $P_2$  which can have at most four finite common zeros, hence

$P_1$ ,  $P_2$  and  $P_3$  can have at most eight finite common zeros in this case.

Case (ii): We will show in this instance that the eleven nodes must be a subset of twelve points of the form  $(\pm\alpha, 0)$ ,  $(0, \pm\beta)$ ,  $(\pm x_1, \pm y_1)$ ,  $(\pm x_2, \pm y_2)$ .

We have both  $\mu_1$  and  $\lambda_1$  nonzero. We first suppose the nodes all lie on

$P^{(2,2)}$ . Now,  $P^{(2,2)}$  and  $P_1$  cannot have a common component; this is easily

established by an argument on the coefficients.  $P^{(2,2)}$  has at most two zeros on each axis, of the form  $(\pm\alpha, 0)$  and  $(0, \pm\beta)$ , as claimed. If  $c_2 = 0$ , the

zeros coalesce at the origin, but we will no longer have enough common zeros.

$Q_2$  and  $P^{(2,2)}$  are seen to have at most eight common zeros, symmetrically located with respect to the axis, i.e.,  $(\pm x_1, \pm y_1)$ ,  $(\pm x_2, \pm y_2)$ .

Now, suppose the nodes do not all lie on  $P^{(2,2)}$ . Then they lie on each of  $P_2 = \mu_2 P^{(4,0)} + \lambda_2 P^{(2,2)}$ , where  $\mu_2 \neq 0$  and  $P_3 = \mu_3 P^{(0,4)} + \lambda_3 P^{(2,2)}$ ,

where  $\mu_3 \neq 0$ . We see that  $P_2$  and  $P_3$  can have at most four common zeros

on the axis. We consider  $Q_2$ ,  $P_2$ , and  $P_3$ . If the three have a common com-

ponent, we see by symmetry that  $Q_2$  must be a factor of both  $P_2$  and  $P_3$ .

Then

$$\begin{aligned} P_2 &= (ax^2 + b)Q_2 \\ P_3 &= (cy^2 + d)Q_2 \end{aligned}$$

Now,  $P_2$  and  $P_3$  have  $(\pm\alpha, 0)$  and  $(0, \pm\beta)$  as common zeros, so  $bQ_2(0, \pm\beta) = 0$  and  $dQ_2(\pm\alpha, 0) = 0$ . If  $b = 0$ ,  $P_2$  has only the origin as a zero on the axis.

This would not be enough zeros. Likewise if  $d = 0$ ,  $P_3$  has only the origin as a zero on the axis, which is impossible. Thus,  $(\pm\alpha, 0)$  and  $(0, \pm\beta)$  lie on  $Q_2$ .

But then all the nodes lie on  $Q_2$ , which is impossible. Thus  $Q_2$  cannot be a factor of both  $P_2$  and  $P_3$ . With no loss of generality, assume  $Q_2$  is not a



factor of  $P_2$ . Then  $Q_2$  and  $P_2$  can have at most 8 common zeros, and they are of the stated form.

It now remains to be shown that no proper subset of twelve points of the form  $(\pm\alpha, 0)$ ,  $(0, \pm\beta)$ ,  $(\pm x_1, \pm y_1)$ ,  $(\pm x_2, \pm y_2)$  can be used for a cubature formula of precision seven for the regions under consideration. Let  $A^{(\alpha, 0)}$  be the weight associated with the node  $(\alpha, 0)$ . Consider  $Q^{(\alpha, 0)} = Q_2 \cdot x \cdot (x + \alpha)$ .

Now,  $Q^{(\alpha, 0)}$  is zero at all the nodes except  $(\alpha, 0)$ , so  $\int_R wQ^{(\alpha, 0)} = A^{(\alpha, 0)} Q^{(\alpha, 0)}_{(\alpha, 0)}$ . Similarly, we have  $\int_R wQ^{(-\alpha, 0)} = A^{(-\alpha, 0)} Q^{(-\alpha, 0)}_{(-\alpha, 0)}$ .

Now we note that because of symmetry  $\int_R wQ^{(\alpha, 0)} = \int_R 2w\alpha x^2$ ,  $\int_R wQ^{(-\alpha, 0)} = -\int_R 2w\alpha x^2$ , and  $Q^{(\alpha, 0)}_{(\alpha, 0)} = -Q^{(-\alpha, 0)}_{(-\alpha, 0)}$ . Thus we have  $A^{(\alpha, 0)} = A^{(-\alpha, 0)}$

and since a ten point formula cannot exist  $A^{(\alpha, 0)} \neq 0$ . Similar consideration of  $Q^{(0, \pm\beta)} = Q_2 \cdot y \cdot (y \mp \beta)^2$  yields  $A^{(0, \beta)} = A^{(0, -\beta)} \neq 0$ .

Now consider

$Q^{(x_i, y_i)} = (x^2 - x_j^2 + y^2 - y_j^2) \cdot x \cdot y \cdot (x + x_i)(y + y_i)$ , where  $j = 3 - i$ ,

for  $i = 1, 2$ . It is easily seen that  $Q^{(x_i, y_i)}$  is zero at all nodes except

$(x_i, y_i)$ . Let  $A^{(x_i, y_i)}$  be the weight associated with  $(x_i, y_i)$ , then we have  $\int_R wQ^{(x_i, y_i)} = A^{(x_i, y_i)} Q^{(x_i, y_i)}_{(x_i, y_i)}$ . Then we note that

$$Q^{(-x_i, y_i)}_{(-x_i, y_i)} = Q^{(x_i, -y_i)}_{(x_i, -y_i)} = Q^{(-x_i, -y_i)}_{(-x_i, -y_i)} = Q^{(x_i, y_i)}_{(x_i, y_i)}$$

and that  $\int_R wQ^{(\pm x_i, \pm y_i)} = \int_R w(x^2 - x_j^2 + y^2 - y_j^2)x^2y^2$ , and thus

$A^{(x_i, y_i)} = A^{(-x_i, y_i)} = A^{(x_i, -y_i)} = A^{(-x_i, -y_i)} \neq 0$ . Thus we have shown that

all 12 points have nonzero weights, and thus no 11 point formula can exist. This concludes the proof of the theorem.

### 3.0 Conclusions

We have shown that 12 is a lower bound for the number of nodes in a cubature formula of precision seven for a symmetric planar region. As noted previously the existence of a 12 point formula for an arbitrary symmetric region has not been shown. To show that the author's construction [1] does not fail it is necessary to show that certain determinants involving the moments are not zero. This problem has been quite intractable, and it seems unlikely the approach of the author can be made to work. Some of the key inequalities used in [1] for fully symmetric regions fail to hold for arbitrary symmetric regions. A considerable effort has been expended in attempting to find a region for which the construction fails, but none has been found. As before the author believes the construction cannot fail, but is unable to prove it.

## REFERENCES

1. Richard Franke, "Minimal Point Cubatures of Precision Seven for Symmetric Planar Regions", SIAM J. Numer. Anal. 10(1973) 849-862.
2. A. H. Stroud, "Integration Formulas and Orthogonal Polynomials II", SIAM J. Numer. Anal. 7(1970) 271-276.
3. A. H. Stroud, Approximate Calculation of Multiple Integrals, Prentice-Hall, Englewood Cliffs, N. J., 1971.

Distribution List

	No. of Copies
Defense Documentation Center Cameron Station Alexandria, Virginia 22314	12
Library Naval Postgraduate School Monterey, California 93940	2
Dean of Research Administration Naval Postgraduate School Monterey, California 93940	2
Professor Richard Franke Department of Mathematics Naval Postgraduate School Monterey, California 93940	5
Professor Craig Comstock Department of Mathematics Naval Postgraduate School Monterey, California 93940	1
Dr. Richard Lau Office of Naval Research Pasadena, California	1
Professor Ladis D. Kovach Acting Chairman, Department of Mathematics Naval Postgraduate School Monterey, California 93940	1
Professor R. E. Barnhill Department of Mathematics University of Utah Salt Lake City, Utah 84112	1
Professor A. H. Stroud Department of Mathematics Texas A & M University College Station, Texas 77843	1
Chief of Naval Research Attn: Mathematics Program Arlington, Virginia 22217	2



U162030

DUDLEY KNOX LIBRARY - RESEARCH REPORTS



5 6853 01067788 3

~~U162030~~