



Calhoun: The NPS Institutional Archive
DSpace Repository

Faculty and Researchers

Faculty and Researchers Collection

2003-09-05

Prediction-skill variability in atmospheric and oceanic models

Chu, Peter C.; Ivanov, Leonid, M.

International Conference on Earth System Modeling: Development and Evaluation of Comprehensive Earth System Models; Atmospheres, Oceans and Sea Ices

<http://hdl.handle.net/10945/36781>

Downloaded from NPS Archive: Calhoun



Calhoun is a project of the Dudley Knox Library at NPS, furthering the precepts and goals of open government and government transparency. All information contained herein has been approved for release by the NPS Public Affairs Officer.

Dudley Knox Library / Naval Postgraduate School
411 Dyer Road / 1 University Circle
Monterey, California USA 93943

<http://www.nps.edu/library>

Prediction-skill variability in atmospheric and oceanic models

Chu, Peter C., and Leonid, M. Ivanov, Naval Postgraduate School, Monterey, California, USA
(chu@nps.navy.mil, tel: 1-831-656-3688)

Various numerical atmospheric and oceanic models have been developed in the past several decades. A fundamental question arises: Can we determine which model provides “the best prediction”? To answer this question, full knowledge of the prediction error statistics of each model is needed. Due to high structural complexity and high dimensionality of the error phase space, establishment of such statistics is difficult. Usually the Gaussian distribution is assumed for the error statistics for simplicity. However, it might not be true for regional ocean models.

A new scalar with the dimension of time, the irreversible-skill time (IT), is defined as the time period when the prediction error first exceeds a pre-determined criterion (i.e., the tolerance level) [originally defined as valid prediction period by Chu et al. (2002b, c)], is introduced to estimate the model predictability for linear and nonlinear stages in the prediction error evolution. The probability density function (PDF) of IT satisfies the backward Fokker-Planck equation (or called Pontryagin-Kolmogorov equation in the Russian literature).

Great advantages of IT for model evaluation are: (1) establishing the analytical framework for estimating the local prediction-skill of regional ocean models, (2) understanding the temporal intermittency of prediction-skill, and (3) searching physical mechanisms causing extremely successful prediction (ESP). Application to ocean (atmospheric) model evaluation is demonstrated.

References

Chu, P.C., L.M. Ivanov and C. Fan, Backward Fokker-Plank equation for determining model valid prediction period. *J. Geophys. Res.*, 107, 6, 11-1 – 11-10, 2002a.

Chu, P.C., Ivanov L.M., Margolina T.M. and O.V. Melnichenko, Probabilistic stability of a simple atmospheric model to differently scales perturbations. *J. Atmos. Sci.*, 59,19,2860-2873, 2002b.

Chu, P.C., L.M. Ivanov, L. Kantha, O.V. Melnichenko ., and Yu. A. Poberezhny, Power decay law in model forecast skill. *Geophys. Res. Let.*, 29 (15), 31-4 -31-8, 2002c.