



Calhoun: The NPS Institutional Archive
DSpace Repository

Faculty and Researchers

Faculty and Researchers' Publications

2003

Flow decomposition for velocity data assimilation

Chu, Peter C.; Ivanov, L.M.; Margolina, T.M.; Melnichenko, O.V.

Chu, P.C., L.M. Ivanov, T.M. Margolina, O. V. Melnichenko, Flow decomposition for velocity data assimilation. Twenty Third General Assembly of the International Union of Geodesy and Geophysics, Sapporo, Japan, 30 June - 11 July 2003.

<http://hdl.handle.net/10945/42401>

This publication is a work of the U.S. Government as defined in Title 17, United States Code, Section 101. Copyright protection is not available for this work in the United States.

Downloaded from NPS Archive: Calhoun



Calhoun is the Naval Postgraduate School's public access digital repository for research materials and institutional publications created by the NPS community. Calhoun is named for Professor of Mathematics Guy K. Calhoun, NPS's first appointed -- and published -- scholarly author.

Dudley Knox Library / Naval Postgraduate School
411 Dyer Road / 1 University Circle
Monterey, California USA 93943

<http://www.nps.edu/library>

FLOW DECOMPOSITION FOR VELOCITY DATA ASSIMILATION

Peter C. Chu, and Leonid M. Ivanov

Naval Postgraduate School, Monterey, California

Tatiana M. Margolina, and Oleg V. Melnichenko

Marine Hydrophysical Institute, National Ukrainian Academy of Sciences
Sevastopol, Crimea

Ocean observational current data are usually acquired from limited number of stations in domains with open boundaries and contain various errors or noises. It is an important task for physical oceanographers to establish (or to reconstruct) a realistic and complete velocity field from sparse and noisy data.

From a mathematical point of view, the reconstruction requires solving a least square problem without or with a priori information (limit) on the circulation characteristics. An a priori limit can be formulated as a set of inequalities that the solutions should satisfy, as a dynamical model applied to the description of circulation dynamics or hypotheses on statistical properties of reconstructed field.

Without knowing statistical weights and without using ocean numerical models, a kinematical method is proposed for reconstructing a velocity field from noisy and sparse data. For a three-dimensional incompressible flow, two scalar functions, toroidal and poloidal potentials, satisfy Poisson equations with the vertical vorticity and vertical velocity as the sources terms, respectively.

In this talk, a new set of basis functions is introduced for reconstructing the ocean circulation in a domain with open boundaries. These functions are the eigenfunctions of Laplacian operator with homogeneous mixed conditions. With known velocities along the open boundary, the mixed boundary conditions are accurate. With unknown velocities along the open boundary, a parameterization scheme is proposed for obtaining approximate open boundary conditions from data. In general, the reconstruction is reduced to linear and nonlinear regression models for known and unknown velocities along the open boundary, respectively. For the latter (without data on the open boundary), the velocity inside the domain and along the boundaries are simultaneously determined.

References

Chu, P.C., L.M. Ivanov, T.P. Korzhova, T.M. Margolina, and O.V. Melnichenko, 2003: Analysis of sparse and noisy ocean current data using flow decomposition. Part 1: Theory, *Journal of Atmospheric and Oceanic Technology*, in press.

Chu, P.C., L.M. Ivanov, T.P. Korzhova, T.M. Margolina, and O.V. Melnichenko, 2003: Analysis of sparse and noisy ocean current data using flow decomposition. Part 2: Applications to Eulerian and Lagrangian Data, *Journal of Atmospheric and Oceanic Technology*, in press.