



**Calhoun: The NPS Institutional Archive**  
**DSpace Repository**

---

Faculty and Researchers

Faculty and Researchers' Publications

---

2017

## Cubic Maiorana-McFarland bent functions with no affine derivative

Mandal, Bimal; Gangopadhyay, Sugata; Stnic, Pantelimon

Taylor & Francis

---

Mandal, Bimal, Sugata Gangopadhyay, and Pantelimon Stnic. "Cubic MaioranaMcFarland bent functions with no affine derivative." *International Journal of Computer Mathematics: Computer Systems Theory* 2.1 (2017): 14-27  
<https://hdl.handle.net/10945/59304>

---

This publication is a work of the U.S. Government as defined in Title 17, United States Code, Section 101. Copyright protection is not available for this work in the United States.

*Downloaded from NPS Archive: Calhoun*



Calhoun is the Naval Postgraduate School's public access digital repository for research materials and institutional publications created by the NPS community. Calhoun is named for Professor of Mathematics Guy K. Calhoun, NPS's first appointed -- and published -- scholarly author.

**Dudley Knox Library / Naval Postgraduate School**  
**411 Dyer Road / 1 University Circle**  
**Monterey, California USA 93943**

<http://www.nps.edu/library>



# Cubic Maiorana–McFarland bent functions with no affine derivative

Bimal Mandal<sup>a</sup>, Sugata Gangopadhyay<sup>b</sup> and Pantellmon Stănică<sup>c</sup>

<sup>a</sup>Department of Mathematics Indian Institute of Technology Roorkee, Roorkee India; <sup>b</sup>Department of Computer Science and Engineering, Indian Institute of Technology Roorkee, Roorkee India; <sup>c</sup>Department of Applied Mathematics, Naval Postgraduate School, Monterey, CA, USA

## ABSTRACT

A class of cubic Maiorana–McFarland ( $\mathcal{M}$ ) bent functions having no affine derivative was constructed by Canteaut and Charpin [*Decomposing bent functions*, IEEE Trans. Inform. Theory 49(8) (2003), pp. 2004–2019], thereby solving an open problem posed by Hou [*Cubic bent functions*, Discrete Math. 189 (1998), pp. 149–161]. The goal of the paper is twofold: we construct two classes of cubic  $\mathcal{M}$  bent functions with no affine derivative and show their mutual affine inequivalence.

## ARTICLE HISTORY

Received 15 July 2016  
Revised 22 January 2017  
Accepted 20 February 2017

## KEYWORDS

Boolean functions; bent functions; differentials; vector spaces; counting

## 1. Introduction

Let  $\mathbb{Z}$  be the ring of integers,  $\mathbb{F}_{2^n}$  be the extension field of degree  $n$  over  $\mathbb{F}_2$ , the prime field of characteristic 2,  $\mathbb{F}_{2^n}^* = \mathbb{F}_{2^n} \setminus \{0\}$ , and for  $m, t \in \mathbb{Z}^+$  (positive integers), let  $\mathbb{F}_{2^t}^m = \{(x_1, x_2, \dots, x_m) : x_i \in \mathbb{F}_{2^t}, 1 \leq i \leq m\}$  be the vector space of dimension  $m$  over  $\mathbb{F}_{2^t}$ . It is customary for the extension field  $\mathbb{F}_{2^{mt}}$  to be identified with  $\mathbb{F}_{2^t}^m$  as  $\mathbb{F}_{2^t}$ -vector spaces and used, interchangeably, where the context allows it. A function from  $\mathbb{F}_{2^n}$  to  $\mathbb{F}_2$  is said to be a *Boolean function* in  $n$  variables. The set of all Boolean functions in  $n$  variables is denoted by  $\mathcal{B}_n$ . The univariate representation of any function  $f \in \mathcal{B}_n$  is

$$f(x) = \sum_{j \in \Gamma(n)} \text{Tr}_1^{n_j}(\alpha_j x^j) + \varepsilon(1 + x^{2^n-1}), \quad (1)$$

where  $\Gamma(n)$  is the set of cyclotomic coset leaders modulo  $2^n - 1$ ,  $n_j$  is the size of the cyclotomic class containing  $j$ ,  $\alpha_j \in \mathbb{F}_{2^{n_j}}$ ,  $\varepsilon = \sum_{x \in \mathbb{F}_{2^n}} f(x) \pmod{2}$ , and  $\text{Tr}_1^k(x) = x + x^2 + x^{2^2} + \dots + x^{2^{k-1}}$ ,  $k \in \mathbb{Z}^+$ , is the trace function. For every  $j \in \mathbb{Z}/(2^n - 1)\mathbb{Z}$ , we can write  $j = \sum_{s \in E} 2^s$  where  $E \subseteq \{0, 1, \dots, n-1\}$ . The cardinality of  $E$ , denoted by  $|E|$ , is referred to as the 2-weight of  $j$  and written as  $w_2(j)$ . The support of  $f \in \mathcal{B}_n$  is  $\text{supp}(f) = \{x \in \mathbb{F}_{2^n} : f(x) \neq 0\}$ , whose cardinality  $\text{wt}(f) = |\text{supp}(f)|$  is the (Hamming) weight of  $f$ . The (Hamming) distance between any two Boolean functions  $f, g \in \mathcal{B}_n$ , denoted by  $d(f, g)$ , is the number of input values at which the outputs of  $f$  and  $g$  disagree, i.e. the cardinality of  $\{x : f(x) \neq g(x), x \in \mathbb{F}_{2^n}\}$ .

The algebraic degree of  $f$  is defined by  $\deg(f) = \max_{j \in \Gamma(n)} \{w_2(j) : \alpha_j \neq 0\}$ . Boolean functions with algebraic degree at most 1 are said to be *affine functions*. Precisely, an affine function  $\varphi_{a,\varepsilon} : \mathbb{F}_{2^n} \rightarrow \mathbb{F}_2$  is of the form

$$\varphi_{a,\varepsilon}(x) = \text{Tr}_1^n(ax) + \varepsilon, \quad \text{for all } x \in \mathbb{F}_{2^n},$$

where  $a \in \mathbb{F}_{2^n}$ ,  $\varepsilon \in \mathbb{F}_2$  (if  $\varepsilon = 0$ , then  $\varphi_{a,0}$  is a linear function). Suppose  $n = mt$ , where  $m, t \in \mathbb{Z}^+$ , and  $f : \mathbb{F}_{2^t}^m \rightarrow \mathbb{F}_2$ . Then the  $m$ -variate representation of  $f$  is

$$f(x_1, \dots, x_m) = \text{Tr}_1^t(g(x_1, \dots, x_m)), \quad \text{for all } (x_1, \dots, x_m) \in \mathbb{F}_{2^t}^m, \quad (2)$$

where  $g(x_1, \dots, x_m) \in \mathbb{F}_{2^t}[x_1, \dots, x_m]$ . In this paper we will mostly use bivariate representations of Boolean functions.

The nonlinearity of  $f \in \mathcal{B}_n$  is  $nl(f) = \min\{d(f, \varphi_{a,\varepsilon}) : a \in \mathbb{F}_{2^n}, \varepsilon \in \mathbb{F}_2\}$ . The Walsh-Hadamard transform of  $f$  at  $a \in \mathbb{F}_{2^n}$  is defined by

$$W_f(a) = \sum_{x \in \mathbb{F}_{2^n}} (-1)^{f(x) + \text{Tr}_1^n(ax)}. \quad (3)$$

The multiset  $\{W_f(a) : a \in \mathbb{F}_{2^n}\}$  is said to be the Walsh-Hadamard spectrum of  $f$ . In the bivariate case, where  $f : \mathbb{F}_{2^t}^2 \rightarrow \mathbb{F}_2$ , instead of Equation (3), we have

$$W_f(a, b) = \sum_{(x,y) \in \mathbb{F}_{2^t}^2} (-1)^{f(x,y) + \text{Tr}_1^t(ax) + \text{Tr}_1^t(by)}, \quad (4)$$

for all  $(a, b) \in \mathbb{F}_{2^t}^2$ . It is known that the nonlinearity of  $f \in \mathcal{B}_n$  can be expressed in terms of the Walsh-Hadamard coefficients as

$$nl(f) = 2^{n-1} - \frac{1}{2} \max_{a \in \mathbb{F}_{2^n}} |W_f(a)| \leq 2^{n-1} - 2^{n/2-1}, \quad n \geq 1, \quad (5)$$

(the upper bound is a direct consequence of Parseval's identity  $\sum_{a \in \mathbb{F}_{2^n}} W_f^2(a) = 2^{2n}$ ). It is known [8] that the upper bound is tight if  $n \in \mathbb{Z}^+$  is even, and achieved by bent functions (first studied by Rothaus [15] and Dillon [9]), which are functions  $f$  for which  $nl(f) = 2^{n-1} - 2^{n/2-1}$ . Also,  $f$  is bent if and only if  $W_f(x) = \pm 2^{n/2}$ , for all  $x \in \mathbb{F}_{2^n}$ , if and only if  $\sum_{x \in \mathbb{F}_{2^n}} (-1)^{f(x) + f(x+a)} = 0$ , for all  $a \in \mathbb{F}_{2^n}^*$ .

### 1.1. Maiorana-McFarland bent functions

Suppose  $n = 2t$  where  $t \in \mathbb{Z}^+$ . Any permutation  $\pi : \mathbb{F}_{2^t} \rightarrow \mathbb{F}_{2^t}$  can be represented by a polynomial  $\pi(x) = \sum_{j=0}^{2^t-1} \alpha_j x^j$  where  $\alpha_j \in \mathbb{F}_{2^t}$ , for all  $0 \leq j \leq 2^t - 1$ . The algebraic degree of  $\pi$  is  $\deg(\pi) := \max\{w_2(j) : \alpha_j \neq 0\}$ . Rothaus [15] proved that any function of the form

$$f : \mathbb{F}_{2^t} \times \mathbb{F}_{2^t} \rightarrow \mathbb{F}_2$$

$$f(x, y) = \text{Tr}_1^t(x\pi(y)) + g(y), \quad \text{for all } (x, y) \in \mathbb{F}_{2^t} \times \mathbb{F}_{2^t}, \quad (6)$$

where  $g \in \mathcal{B}_t$ , is bent. These bent functions are said to be Maiorana-McFarland bent functions and their set is denoted by  $\mathcal{M}$ . In this paper we assume  $g$  to be identically zero. For  $f \in \mathcal{M}$  with  $g = 0$  the algebraic degree is  $\deg(f) = \deg(\pi) + 1$ . Maiorana-McFarland construction provides a natural connection between permutations over finite fields and functions in  $\mathcal{M}$ . Permutations having algebraic degree 1 are said to be linearized permutations. Each linearized permutation on  $\mathbb{F}_{2^t}$  generates a quadratic function in  $\mathcal{M}$ .

Suppose that  $\gcd(t, i) = e$ , with  $\frac{t}{e}$  an odd positive integer, and  $\alpha \in \mathbb{F}_{2^t}$  such that  $\alpha \neq \zeta^{m(2^t-1)}$  for any  $m \in \mathbb{Z}$  and any primitive element  $\zeta$  of  $\mathbb{F}_{2^t}$ . Blokhuis *et al.* [1] mention that  $\sigma_j, j \in \{1, 2, 3\}$ , listed

below are linearized permutations on  $\mathbb{F}_{2^t}$ .

$$\begin{aligned}\sigma_1(x) &= x^{2^i} + \alpha x, \\ \sigma_2(x) &= x^{2^{2i}} + \alpha^{2^i} x, \\ \sigma_3(x) &= x^{2^i} + \alpha^{2^i} x.\end{aligned}\tag{7}$$

Some other linearized polynomials over  $\mathbb{F}_{2^t}$  which will be used in the paper are as follows:

$$\begin{aligned}\sigma_4(y) &= y^{2^{2i}} + \alpha^{2^{2i}} y, \\ \sigma_5(y) &= y^{2^{2i}} + \alpha y, \\ \sigma_6(y) &= y + \alpha^{2^i} y^{2^i}, \\ \sigma_7(y) &= y + \alpha^{2^{2i}} y^{2^{2i}}, \\ \sigma_8(y) &= y + \alpha^{2^{2i}} y^{2^i}.\end{aligned}\tag{8}$$

Observe that the linearized function  $\sigma_4(y) = 0$  if and only if  $y = 0$  or  $y^{2^{2i}-1} = \alpha^{2^{2i}}$ . If  $y^{2^{2i}-1} = \alpha^{2^{2i}}$ , then  $(\alpha^{(2^i-1)/(2^e-1)})^{2^{2i}} = 1$ , since  $e \mid i$ , which implies  $\alpha^{(2^i-1)/(2^e-1)} = 1$ . This is a contradiction, since  $\alpha^{(2^i-1)/(2^e-1)} \neq 1$ . Thus,  $\sigma_4$  is a linearized permutation. Similarly, it can be proved that  $\sigma_j, j = 5, 6, 7, 8$  are linearized permutations.

Each function  $f(x, y) = \text{Tr}_1^t(x\sigma_j(y))$ ,  $1 \leq j \leq 8$ , is a quadratic bent in  $\mathcal{M}$ . Moreover, the following two quadratic permutations were constructed by Blockhuis *et al.* [1]:

$$\begin{aligned}\pi_1(y) &= y^{2^i+1} + \alpha y^{2^{t-i}+1}, \\ \pi_2(y) &= y(\text{Tr}_t^t(y) + \alpha y),\end{aligned}\tag{9}$$

where  $t = k\ell$ ,  $k$  is an odd integer and  $\ell > 1$  is any positive integer (discussed later in details in Section 2.1) on the parameter  $\alpha$ . In this paper, we use the functions of the form  $f_j(x, y) = \text{Tr}_1^t(x\pi_j(y))$ ,  $1 \leq j \leq 2$  as a source of cubic bent functions and consider their differential properties.

### 1.2. Preliminary results

Recall the following well known facts from elementary number theory, which we use frequently in this paper. Suppose that  $ax \equiv b \pmod{n}$  where  $a, b, n \in \mathbb{Z}$  and  $d = \text{gcd}(a, n)$ . Then

- (1) if  $d$  does not divide  $b$ , the congruence has no solution;
- (2) if  $d$  divide  $b$  then all solutions of the congruence are  $x_0 + k(n/d)$ ,  $0 \leq k < d$ , where  $x_0$  is the unique solution to  $(a/d)x \equiv (b/d) \pmod{n/d}$ .

Let  $t$  be a positive integer and  $\text{gcd}(t, i) = e$ . Then [7, p. 2]

$$\text{gcd}(2^{2i} - 1, 2^t - 1) = 2^{\text{gcd}(2i, t)} - 1 = \begin{cases} 2^e - 1 & \text{if } \frac{t}{e} \text{ is odd;} \\ 2^{2e} - 1 & \text{if } \frac{t}{e} \text{ is even.} \end{cases}$$

**Theorem 1.1** ([7, Theorem 3.1]). Let  $\zeta$  be a primitive element of  $\mathbb{F}_{2^t}$  and  $\gcd(t, i) = e$ . For any  $a \in \mathbb{F}_{2^t}^*$ , consider the equation  $a^{2^i} x^{2^{2i}} + ax = 0$  over  $\mathbb{F}_{2^t}$ . Then:

- (1) If  $t/e$  is odd then there are  $2^e$  solutions to this equation for any choice of  $a \in \mathbb{F}_{2^t}^*$ .
- (2) If  $t/e$  is even then there are two possible cases:
  - (a) if  $a = \zeta^{s(2^e+1)}$  for some  $s$ , then there are  $2^{2e}$  solutions to the equation.
  - (b) if  $a \neq \zeta^{s(2^e+1)}$  for any  $s$ , then there exists one solution only, namely  $x = 0$ .

### 1.3. Affine equivalence

The general linear group of degree  $n$  over  $\mathbb{F}_2$ , denoted by  $GL(n, \mathbb{F}_2)$ , is the group of invertible linear transformations acting on  $\mathbb{F}_2^n$ . For any  $A \in GL(n, \mathbb{F}_2)$  and  $x \in \mathbb{F}_2^n$  we denote the action of  $A$  on  $x$  by  $x \mapsto xA$ . The affine general linear group,  $AGL(n, \mathbb{F}_2)$ , is the set of all transformations of the form  $x \mapsto xA + b$  where  $b \in \mathbb{F}_2^n$ . This group can be thought of as the semidirect product  $GL(n, \mathbb{F}_2) \ltimes \mathbb{F}_2^n$ , but we will not need that here.

**Definition 1.1.** Two Boolean functions  $f, g \in \mathcal{B}_n$  are said to be affine equivalent if there exists  $(A, b) \in AGL(n, \mathbb{F}_2)$  such that  $g(x) = f(xA + b)$ , for all  $x \in \mathbb{F}_2^n$ .

For Boolean functions used as cryptographic primitives the notion of equivalence is further generalized as follows.

**Definition 1.2.** Two Boolean functions  $f, g \in \mathcal{B}_n$  are said to be extended affine equivalent (EA-equivalent) if there exist  $(A, b) \in AGL(n, \mathbb{F}_2)$ ,  $a \in \mathbb{F}_2^n$  and  $\varepsilon \in \mathbb{F}_2$  such that  $g(x) = f(xA + b) + \varphi_{a,\varepsilon}(x)$ , for all  $x \in \mathbb{F}_2^n$  where  $\varphi_{a,\varepsilon}(x) = \text{Tr}_1^n(ax) + \varepsilon$ .

If two Boolean functions  $f, g \in \mathcal{B}_n$  have different algebraic degrees then they are EA-inequivalent. Therefore, the algebraic degree serves as an EA-invariant. The multiset consisting of absolute values of Walsh–Hadamard transforms of a function  $f$  is said to be its *absolute Walsh–Hadamard spectrum*. If the absolute Walsh–Hadamard spectra of two Boolean functions are different, which is possible even if their algebraic degrees are the same, then we know that they are EA-inequivalent. Thus, the absolute Walsh–Hadamard spectrum serves as a more sophisticated EA-invariant. In fact, the autocorrelation spectrum which is another invariant is also connected to the Walsh–Hadamard spectrum. For bent functions the absolute Walsh–Hadamard spectrum is unique and flat, set to  $2^{n/2}$  where  $n$  is the number of variables. For this reason the invariants dependent on Walsh–Hadamard spectra are unable to decide EA-inequivalence of bent functions. In this paper, we use a second-derivative-based invariant to distinguish between the classes of cubic bent functions in  $\mathcal{M}$  which have no affine derivative.

### 1.4. Derivatives and affine inequivalence

The problem of deciding EA-inequivalence is completely solved for Boolean functions having algebraic degrees at most 2, that is, for affine and quadratic Boolean functions. We refer to MacWilliams and Sloane [14, Chapter 15] for detailed discussion on quadratic Boolean functions including their affine inequivalence. In the absence of a general theory for functions having algebraic degree three and above we address the problem by considering derivatives of these functions.

**Definition 1.3.** The derivative of  $f \in \mathcal{B}_n$  with respect to an  $m$ -dimensional  $\mathbb{F}_2$ -subspace  $V$  of  $\mathbb{F}_2^n$ , or the  $m$ th-(order) derivative, is the function  $D_V f : \mathbb{F}_2^n \rightarrow \mathbb{F}_2$  defined by

$$D_V f(x) = \sum_{a \in V} f(x + a), \quad \text{for all } x \in \mathbb{F}_2^n. \quad (10)$$

The algebraic degree to  $D_V f$  is at most  $\deg(f) - m$ . If  $V$  is one-dimensional then  $D_V f(x) = f(x + a) + f(x)$  where  $a \in V \setminus \{0\}$ , which is usually denoted by  $D_a f(x)$ . If  $V$  is a 2-dimensional subspace of  $\mathbb{F}_{2^n}$  we choose any pair of distinct elements  $a, b \in V \setminus \{0\}$  and write

$$D_V f(x) = D_{a,b} f(x) = f(x) + f(x + a) + f(x + b) + f(x + a + b),$$

for all  $x \in \mathbb{F}_{2^n}$ . Obviously the choice of  $(a, b)$  does not change the function  $D_V f$ .

Dillon [9] proposed proving inequivalence of Boolean functions by considering their  $m$ th-order derivatives over all distinct  $m$ -dimensional subspaces of  $\mathbb{F}_{2^n}$ .

**Theorem 1.2 ([9, Theorem 2.1]).** *For any function  $f \in \mathcal{B}_n$  let  $\mathcal{D}_k(f)$  denote the multiset of all  $k$ -dimensional derivatives of  $f$ . If  $f, g \in \mathcal{B}_n$  are affinely equivalent, then so are  $\mathcal{D}_k(f)$  and  $\mathcal{D}_k(g)$ . If the nonsingular affine transformation  $A$  (operating on  $\mathcal{B}_n$ ) maps  $f$  onto  $g$ , then it also maps  $\mathcal{D}_k(f)$  onto  $\mathcal{D}_k(g)$ .*

Dillon proved the following corollary to Theorem 1.2.

**Corollary 1.1.** *If  $\mathcal{P}$  is any affine invariant for  $\mathcal{B}_n$ , then  $f \rightarrow \mathcal{P}(\mathcal{D}_n(f))$  is also an affine invariant for  $\mathcal{B}_n$ .*

Derivatives have been used for this purpose by Carlet [4] and Canteaut and Charpin [3]. Second derivatives have been used by Gangopadhyay [11] extensively to demonstrate affine inequivalence between cubic bent function in  $\mathcal{M}$  which are in many ways similar to each other. The technique can be summarized as follows:

- (1) For  $f \in \mathcal{B}_n$ , construct the set

$$S_f = \{wt(D_V f) : V \text{ varies over all distinct two-dimensional subspaces of } \mathbb{F}_2^n\}.$$

- (2) Construct the frequency distribution of the weights in  $S_f$ . We refer to  $S_f$  as the second-derivative weight distribution of  $f$ .
- (3) For any two  $f, g \in \mathcal{B}_n$ , if the second-derivative weight distributions of  $f$  and  $g$  are different then  $f$  and  $g$  are affine inequivalent.

### 1.5. A problem proposed by Hou

Hou [12] proposed the problem of finding values of  $n$  for which there exist  $n$ -variable cubic bent functions having no affine derivative. Canteaut and Charpin [3, Lemma 1] considered the cubic bent functions of the form  $f(x, y) = \text{Tr}_1^t(xy^{2^t+1})$  and proved that they have no affine derivative. Using this result they constructed an infinite family of cubic bent functions. Gangopadhyay [11] identified subclasses of inequivalent bent functions within this class, by using an invariant proposed by Dillon [9].

**Theorem 1.3 ([11, Theorem 4]).** *Let  $n = 2t$ . If  $f_i(x, y) = \text{Tr}_1^t(xy^{2^t+1})$  where  $x, y \in \mathbb{F}_{2^t}$ ,  $n \geq 6$ ,  $i \in \mathbb{Z}$  such that  $1 \leq i < t$  and  $\gcd(2^i + 1, 2^t - 1) = 1$ , then the number of constant functions among  $D_V f_i$  is*

$$\frac{(2^t - 1)(2^{t+e-1}(2^e + 1) - (2^t + 1))}{3}$$

where  $\gcd(t, i) = e$ .



Using this result, Gangopadhyay [11, Corollary 5] proved that if  $\gcd(t, i) \neq \gcd(t, j)$  then  $f_i$  and  $f_j$  are not affine equivalent. In this paper our goal is not only to identify some more classes of cubic bent functions in  $\mathcal{M}$  having no affine derivative but also to prove affine inequivalence between the classes of functions so obtained. We use Theorem 1.3 almost exclusively for that purpose.

### 1.6. Motivation

Boolean functions are used in many cryptosystems. For example, in some LFSR-based stream ciphers, a Boolean function is used to combine the outputs of several LFSRs. To design a secure cryptosystem, it is required that the output of the Boolean function should not be correlated with a subset of input variables, and thus, to resist correlation attacks [16], the concept of resiliency of a Boolean function was invented. In [2, Proposition 4.2], Camion et al. constructed a resilient function which is quite similar to the Maiorana–McFarland bent functions. Further, in a block cipher, linear structures have been investigated for their cryptanalytic significance. In [6], Charpin et al. derived a relation between polynomial with linear structure and a Maiorana–McFarland function with an affine derivative.

In the recent past some new cryptographic properties have been proposed which are relevant particularly when functions depending on a small number of variables are used as cryptographic primitives. One such property is the second-order nonlinearity of a Boolean function [5], that is, the Hamming distance of the function from the set of quadratic Boolean functions. The experimental evidence [10, Section 3] suggests that cubic bent functions having no affine derivative might be possessing higher second-order derivatives than the rest.

Thus, Boolean functions that do or do not possess affine derivative are important from a cryptographic perspective. In this paper we show that the functions of the form  $f_j(x, y) = \text{Tr}_1^t(x\pi_j(y))$ ,  $1 \leq j \leq 2$ , where  $\pi_j$ 's are defined by Equation (9), do not have affine derivative and they are mutually affine inequivalent.

Our analysis shows that there exist quite a few affine inequivalent classes of cubic bent function in  $\mathcal{M}$  having no affine derivative. Primary research direction ought to be towards finding out such bent functions outside  $\mathcal{M}$  and evaluating the significance of those functions in cryptography and coding theory.

## 2. Cubic bent functions in $\mathcal{M}$

Two subclasses of cubic bent functions in  $\mathcal{M}$  are constructed by using the permutations in (9). We show that the functions in each of these classes have no affine derivative. We prove that the functions in the different subclasses are affine inequivalent by considering their second-derivative weight distributions. Thus we extend the number of known cubic bent functions in  $\mathcal{M}$  with no affine derivative.

### 2.1. Subclass associated with $\pi_1(y) = y^{2^i+1} + \alpha y^{2^{t-i}+1}$

Let  $n = 2t$ ,  $t \geq 3$  and  $\zeta$  be a primitive element of  $\mathbb{F}_{2^t}$ . Blokhuis et al. [1] proved that the function  $\pi_1: \mathbb{F}_{2^t} \rightarrow \mathbb{F}_{2^t}$  defined by

$$\pi_1(y) = y^{2^i+1} + \alpha y^{2^{t-i}+1},$$

for all  $y \in \mathbb{F}_{2^t}$ , where  $i \in \mathbb{Z}$  such that  $1 \leq i < t$  is a permutation if the following conditions are satisfied:

1.  $\gcd(i, t) = e$  and  $t/e$  is odd;
2.  $\alpha \neq \zeta^{s(2^e-1)}$ , for any  $s \in \mathbb{Z}$ .

**Lemma 2.1.** Under the above conditions, the cubic Maiorana-McFarland bent function  $f_i : \mathbb{F}_{2^t} \times \mathbb{F}_{2^t} \rightarrow \mathbb{F}_2$  defined by

$$f_i(x, y) = \text{Tr}_1^t(xy^{2^i+1} + \alpha xy^{2^{t-i}+1}), \quad \text{for all } (x, y) \in \mathbb{F}_{2^t} \times \mathbb{F}_{2^t}, \quad (11)$$

does not possess any affine derivative.

**Proof:** Let  $a, b \in \mathbb{F}_{2^t}$ . Then the first derivative of  $f_i$  at  $(a, b) \in \mathbb{F}_{2^t} \times \mathbb{F}_{2^t}$  is

$$\begin{aligned} D_{(a,b)}f_i(x, y) &= \text{Tr}_1^t(a\pi_1(y) + (x+a)D_b\pi_1(y)) \\ &= \text{Tr}_1^t(a(y^{2^i+1} + \alpha y^{2^{t-i}+1}) \\ &\quad + (x+a)(y^{2^i}b + yb^{2^i} + b^{2^i+1} + \alpha y^{2^{t-i}}b + \alpha yb^{2^{t-i}} + \alpha b^{2^{t-i}+1})), \end{aligned}$$

for all  $(x, y) \in \mathbb{F}_{2^t} \times \mathbb{F}_{2^t}$ . If  $a \neq 0$ , then  $D_{(a,b)}f_i(x, y)$  is a quadratic function. If  $a = 0$  and  $b \neq 0$ , then

$$D_{(0,b)}f_i(x, y) = \text{Tr}_1^t(x(y^{2^i}b + yb^{2^i} + \alpha y^{2^{t-i}}b + \alpha yb^{2^{t-i}}) + x(b^{2^i+1} + \alpha b^{2^{t-i}+1}))$$

is an affine function if and only if  $p(y) = y^{2^i}b + yb^{2^i} + \alpha y^{2^{t-i}}b + \alpha yb^{2^{t-i}}$  is constant for all  $y \in \mathbb{F}_{2^t}$ . Since  $p(0) = 0$ , then (simplifying by  $y$  above)  $y^{2^i-1} + b^{2^i-1} + \alpha y^{2^{t-i}-1} + \alpha b^{2^{t-i}-1} = 0$ , for all  $y \in \mathbb{F}_{2^t}^*$ . For  $y = 1$ , we get  $b^{2^i-1} + \alpha b^{2^{t-i}-1} = 1 + \alpha$ , which renders

$$y^{2^i-1} + \alpha y^{2^{t-i}-1} + 1 + \alpha = 0. \quad (12)$$

If  $\alpha = 0$ , then the solution space of Equation (12) is  $\mathbb{F}_{2^e}$ . If  $\alpha \neq 0$ , we know that for  $y \in \mathbb{F}_{2^e}$ , then  $y^{2^i-1} = 1 = y^{2^{t-i}-1}$ , since  $e = \gcd(i, t)$ . Therefore Equation (12) is identically zero. Otherwise, substituting  $y = c \in \mathbb{F}_{2^t} \setminus \mathbb{F}_{2^e}$  in Equation (12)  $c^{2^i-1} + \alpha c^{2^{t-i}-1} + 1 + \alpha = 0$ , so  $\alpha = (c + c^{2^i}) / (c + c^{2^{t-i}})$  and then,  $\alpha^{2^i} = (c + c^{2^i})^{2^i} / (c + c^{2^{t-i}})^{2^i} = (c + c^{2^i})^{2^i-1}$ , that is,  $(\alpha^{(2^i-1)/(2^e-1)})^{2^i} = 1$ , since  $e \mid i$ , which implies  $\alpha^{(2^i-1)/(2^e-1)} = 1$ . This is a contradiction, since  $\alpha^{(2^i-1)/(2^e-1)} \neq 1$  (otherwise, the condition  $\alpha \neq \zeta^{s(2^e-1)}$  would be violated). Thus Equation (12) does not hold for all  $y \in \mathbb{F}_{2^t}$ . Therefore,  $D_{(0,b)}f_i$  is not an affine function, and our lemma is shown. ■

**Theorem 2.1.** The number of distinct two-dimensional subspaces corresponding to constant second-derivatives of  $f_i$  is

$$\frac{(2^t - 1)(2^{t+e-1}(2^e + 1) - (2^t + 1))}{3}.$$

**Proof:** Let  $V = \langle (a, b), (c, d) \rangle$  be any 2-dimensional subspace of  $\mathbb{F}_{2^t} \times \mathbb{F}_{2^t}$ . The second order derivative of  $f_i$  is

$$\begin{aligned} D_V f_i(x, y) &= \text{Tr}_1^t((ad + bc)y^{2^i} + (ad^{2^i} + cb^{2^i})y + \alpha(ad + cb)y^{2^{t-i}} + \alpha(ad^{2^{t-i}} + cb^{2^{t-i}})y \\ &\quad + \gamma x + (ad^{2^i+1} + cb^{2^i+1}) + \alpha(ad^{2^{t-i}+1} + cb^{2^{t-i}+1}) + (a + c)\gamma), \end{aligned}$$

where  $\gamma = (bd^{2^i} + b^{2^i}d) + \alpha(bd^{2^{t-i}} + b^{2^{t-i}}d)$ .

*Case 1:* We first assume  $b = 0, d = 0$ . Then  $D_V f_i(x, y) = 0$  for all  $(x, y) \in \mathbb{F}_{2^t} \times \mathbb{F}_{2^t}$ . Thus, with respect to any 2-dimensional subspace of  $\mathbb{F}_{2^t} \times \{0\}$  the second order derivative of  $f_i$  is 0. Therefore the number of two-dimensional subspaces such that  $D_V f_i$  is constant, is equal to  $(2^t - 1)(2^{t-1} - 1)/3$ .



Case 2: Let  $b = 0$  but  $d \neq 0$ . Then

$$\begin{aligned} D_V f_i(x, y) &= \text{Tr}_1^t((ad)y^{2^i} + (ad^{2^i})y + \alpha(ad)y^{2^{i-1}} + \alpha(ad^{2^{i-1}})y + (ad^{2^i+1}) + \alpha(ad^{2^{i-1}+1})) \\ &= \text{Tr}_1^t(((ad)^{2^{i-1}} + ad^{2^i} + (\alpha ad)^{2^i} + \alpha ad^{2^{i-1}})y) + \text{Tr}_1^t(ad^{2^i+1} + \alpha ad^{2^{i-1}+1}). \end{aligned}$$

Thus,  $D_V f_i(x, y)$  is constant if and only if  $(ad)^{2^{i-1}} + ad^{2^i} + (\alpha ad)^{2^i} + \alpha ad^{2^{i-1}} = 0$ , and so,

$$\begin{aligned} 0 &= ad + a^{2^i} d^{2^{2i}} + (\alpha ad)^{2^{2i}} + (\alpha a)^{2^i} d \\ &= (a^{2^i} + (\alpha a)^{2^{2i}}) d^{2^{2i}} + (a + (\alpha a)^{2^i}) d, \end{aligned}$$

which can be written as  $h^{2^i} d^{2^{2i}} + hd = 0$ , where  $h = a + (\alpha a)^{2^i}$ . Thus,  $h \neq 0$  as  $a \neq 0$ . Then, by Theorem 1.1, the above equation has  $2^e - 1$  nonzero solutions for  $d$  in  $\mathbb{F}_{2^t}$ . Therefore, for any nonzero  $a \in \mathbb{F}_{2^t}$ , it is possible to choose  $d$  in  $2^e - 1$  ways,  $a$  can be chosen in  $2^t - 1$  ways and  $c$  in  $2^t$  ways, since the subspace generated by  $\{(a, 0), (c, d)\}$  is equal to the subspace generated by  $\{(a, 0), (a + c, d)\}$ . Therefore, the total number of two-dimensional subspaces such that the second derivative of  $f_i$  is constant, is equal to  $(2^t - 1)2^{t-1}(2^e - 1)$ .

Case 3: Let  $b \neq 0$  and  $d \neq 0$ .

Subcase (i): Let  $b = d$ . Then the subspace generated by  $\{(a, b), (c, d)\}$  is equal to the subspace generated by  $\{(a + c, b + d), (c, d)\} = \{(a + c, 0), (c, d)\}$ , which is the same as in the previous case.

Subcase (ii): Let  $b \neq d$ . Then,  $D_V f_i(x, y)$  is constant if and only if

$$\text{Tr}_1^t((ad + bc)y^{2^i} + (ad^{2^i} + cb^{2^i})y + \alpha(ad + cb)y^{2^{i-1}} + \alpha(ad^{2^{i-1}} + cb^{2^{i-1}})y) = 0 \quad (13)$$

and

$$\gamma = 0. \quad (14)$$

From (14), we have  $b^{2^i-1} + \alpha b^{2^{i-1}-1} = d^{2^i-1} + \alpha d^{2^{i-1}-1}$ , since  $b \neq 0$  and  $d \neq 0$ . Again, from Equation (14),

$$0 = \gamma = \gamma^{2^i} = b^{2^i} d^{2^{2i}} + b^{2^{2i}} d^{2^i} + \alpha^{2^i} (b^{2^i} d + b d^{2^i}) = (bd^{2^i} + b^{2^i} d)^{2^i} + \alpha^{2^i} (b^{2^i} d + b d^{2^i}).$$

Let  $z = bd^{2^i} + b^{2^i} d$ . Then the above equation can be written as  $z^{2^i} + \alpha^{2^i} z = 0$ , which has the only solution  $z = 0$ , that is,

$$b^{2^i} d + b d^{2^i} = 0, \iff \left(\frac{d}{b}\right)^{2^i-1} = 1, \quad \text{as } b \neq 0 \text{ and } d \neq 0, \quad (15)$$

$$\text{and so, } \frac{d}{b} \in \mathbb{F}_{2^e}^*, \quad \text{as } \gcd(t, i) = e.$$

Since  $b \neq d$ , for any nonzero  $b$ , there exist a nonzero  $\lambda \in \mathbb{F}_{2^e}$  with  $\lambda \neq 1$  such that  $d = \lambda b$ . Thus,  $d$  can be chosen in  $2^e - 2$  ways and  $b$  in  $2^t - 1$  ways.

Further, from Equation (13), we have

$$\begin{aligned} \text{Tr}_1^t((ad + bc)y^{2^i} + \alpha(ad + cb)y^{2^{i-1}} + y((ad^{2^i} + cb^{2^i}) + \alpha(ad^{2^{i-1}} + cb^{2^{i-1}}))) &= 0, \\ \iff \text{Tr}_1^t((ad + bc)y^{2^i} + \alpha(ad + cb)y^{2^{i-1}} + y(ad + bc)(b^{2^i-1} + \alpha b^{2^{i-1}-1})) &= 0, \\ \iff \text{Tr}_1^t((ad + bc)^{2^{i-1}} + (\alpha(ad + cb))^{2^i} + (ad + bc)(b^{2^i-1} + \alpha b^{2^{i-1}-1}))y) &= 0, \end{aligned}$$

for all  $y \in \mathbb{F}_{2^t}$  if and only if the following (equivalent) statements hold:

$$\begin{aligned} (ad + bc)^{2^{t-1}} + (\alpha(ad + cb))^{2^t} + (ad + bc)(b^{2^t-1} + \alpha b^{2^{t-1}-1}) &= 0, \\ \Leftrightarrow b^{2^{t-1}}(a\lambda + c)^{2^{t-1}} + b^{2^t}(\alpha(a\lambda + c))^{2^t} + (a\lambda + c)(b^{2^t} + \alpha b^{2^{t-1}}) &= 0, \\ \Leftrightarrow b^{2^{t-1}}w^{2^{t-1}} + b^{2^t}(\alpha w)^{2^t} + w(b^{2^t} + \alpha b^{2^{t-1}}) &= 0, \quad \text{where } w = a\lambda + c, \\ \Leftrightarrow bw + (\alpha b)^{2^{2t}}w^{2^{2t}} + (b^{2^{2t}} + \alpha^{2^t}b)w^{2^t} &= 0, \\ \Leftrightarrow w((\alpha b)^{2^{2t}}w^{2^{2t-1}} + (b^{2^{2t}} + \alpha^{2^t}b)w^{2^t-1} + b) &= 0. \end{aligned}$$

Therefore, we infer that either  $w=0$  or  $(\alpha b)^{2^{2t}}w^{2^{2t-1}} + (b^{2^{2t}} + \alpha^{2^t}b)w^{2^t-1} + b = 0$ , which can be transformed into

$$\begin{aligned} (\alpha b)^{2^{2t}}w^{(2^t-1)(2^t+1)} + (b^{2^{2t}} + \alpha^{2^t}b)w^{2^t-1} + b &= 0, \\ \Leftrightarrow \alpha^{2^{2t}}b^{2^{2t}}\mu^{2^t+1} + b^{2^{2t}}\mu + \alpha^{2^t}b\mu + b &= 0, \quad \text{where } w^{2^t-1} = \mu, \\ \Leftrightarrow (\alpha^{2^t}\mu + 1)^{2^t}b^{2^{2t}}\mu + (\alpha^{2^t}\mu + 1)b &= 0, \\ \Leftrightarrow b(\alpha^{2^t}\mu + 1)(b^{2^{2t}-1}\mu(\alpha^{2^t}\mu + 1)^{2^t-1} + 1) &= 0, \\ \alpha^{2^t}\mu + 1 \neq 0, \text{ since the only solution of } \alpha^{2^t}w^{2^t} + w = 0 &\text{ is } w = 0 \text{ due to } \sigma_6(y), \\ \Leftrightarrow b^{2^{2t}-1}\mu(\alpha^{2^t}\mu + 1)^{2^t-1} + 1 = 0, \text{ as } b \neq 0 \text{ and } \alpha^{2^t}\mu + 1 \neq 0, \\ \Leftrightarrow b^{(2^t+1)(2^t-1)}w^{2^t-1}(\alpha^{2^t}w^{2^t-1} + 1)^{2^t-1} + 1 &= 0, \\ \Leftrightarrow (b^{2^t+1}(\alpha^{2^t}w^{2^t} + w))^{2^t-1} = 1, \\ \Leftrightarrow b^{2^t+1}(\alpha^{2^t}w^{2^t} + w) \in \mathbb{F}_{2^t}^*, \\ \Leftrightarrow b^{2^t+1}(\alpha^{2^t}w^{2^t} + w) \in \mathbb{F}_{2^e}^*, \text{ as } \gcd(i, t) = e, \end{aligned}$$

and thus

$$\alpha^{2^t}w^{2^t} + w = \frac{\lambda'}{b^{2^t+1}}, \quad \text{as } b \neq 0 \text{ and } \lambda' \in \mathbb{F}_{2^e}. \quad (16)$$

Since the homogeneous part of the above equation is a linear equation which has a unique solution  $w=0$ , then Equation (16) has a unique solution in  $\mathbb{F}_{2^t}$  for each  $\lambda' \in \mathbb{F}_{2^e}$ . Thus  $w$  can be chosen in  $2^e$  ways (including  $w=0$ ). For fixed  $a$  and  $b$ ,  $c$  can be chosen in  $2^e$  ways. Therefore,  $a$  can be chosen in  $2^t$  ways,  $b$  in  $2^t - 1$  ways,  $d$  in  $2^e - 2$  ways and  $c$  in  $2^e$  ways. Each 2-dimensional subspace generated by a pair of vectors  $(a, b)$  and  $(c, d)$  satisfying the above conditions, contains altogether 6 distinct bases satisfying these conditions. Therefore, the total number of distinct two-dimensional subspaces with bases of this type is  $2^{t+e}(2^t - 1)(2^e - 2)/6$ . Adding the counts from the above three cases we obtain the total count  $(2^t - 1)(2^{t+e-1}(2^e + 1) - (2^t + 1))/3$ , and the theorem is shown.  $\blacksquare$

**Remark 2.1:** If  $\alpha = 0$ , then the cubic Maiorana-McFarland bent function, defined as in Equation (11) is  $f_i(x, y) = \text{Tr}_1^t(xy^{2^t+1})$ , for all  $(x, y) \in \mathbb{F}_{2^t} \times \mathbb{F}_{2^t}$ . From Theorem 1.3, we have the number of constant functions among the second order derivative of  $f_i$  is  $(2^t - 1)(2^{t+e-1}(2^e + 1) - (2^t + 1))/3$ .

## 2.2. The subclass associated with $\pi_2(y) = y(\text{Tr}_t^e(y) + \alpha y)$

We next consider a class of permutation polynomials constructed by Blokhuis [1] and referred to by Laigle-Chapuy [13].

**Theorem 2.2** ([1], [13, Theorem 2]). Let  $t = k\ell$ , where  $k$  be an odd and  $\ell > 1$  be any positive integer. Then the following polynomial is a bilinear permutation over  $\mathbb{F}_{2^t}$  of the form:

$$\pi(x) = x(\text{Tr}_\ell^t(x) + \alpha x),$$

where  $\alpha \in \mathbb{F}_{2^t} \setminus \mathbb{F}_2$  and  $\text{Tr}_\ell^t(x) = \sum_{i=0}^{k-1} x^{2^{i\ell}}$ .

Using this class of permutations we construct a class of cubic Maiorana–McFarland bent functions. Let  $t = k\ell$ , where  $k$  be an odd and  $\ell > 1$  be any positive integer. A function  $g : \mathbb{F}_{2^t} \times \mathbb{F}_{2^t} \rightarrow \mathbb{F}_2$  defined by

$$g(x, y) = \text{Tr}_1^t(xy \text{Tr}_\ell^t(y) + \alpha xy^2), \quad \text{for all } (x, y) \in \mathbb{F}_{2^t} \times \mathbb{F}_{2^t}, \quad (17)$$

is a cubic Maiorana–McFarland bent. We prove that if  $k > 1$ , then the functions  $g$  belonging to this class do not have any affine derivative.

**Theorem 2.3.** Let  $t = k\ell$ , where  $k$  be an odd and  $\ell > 1$  be any positive integer. If  $k > 1$ , then the cubic Maiorana–McFarland bent function  $g$  defined as in Equation (17) has no affine derivative.

**Proof:** Let  $(a, b)$  be an any element of  $\mathbb{F}_{2^t} \times \mathbb{F}_{2^t}$ .

$$\begin{aligned} D_{(a,b)}g(x, y) &= g(x, y) + g(x + a, y + b) \\ &= \text{Tr}_1^t(xy \text{Tr}_\ell^t(y) + (x + a)(y + b) \text{Tr}_\ell^t(y + b) + \alpha xy^2 + \alpha(x + a)(y + b)^2) \\ &= \text{Tr}_1^t(a(y \text{Tr}_\ell^t(y) + \alpha y^2) + (x + a)(y \text{Tr}_\ell^t(b) + b \text{Tr}_\ell^t(y) + b \text{Tr}_\ell^t(b) + \alpha b^2)). \end{aligned}$$

Let  $a \neq 0$ . Since  $y \text{Tr}_\ell^t(y) + \alpha y^2 = 0 \iff y = 0$  or  $\text{Tr}_\ell^t(y) = \alpha y \iff y = 0$ . Thus, if  $a \neq 0$ ,  $D_{(a,b)}g$  is a quadratic function. Let us consider  $a = 0$ , so

$$D_{(0,b)}g(x, y) = \text{Tr}_1^t(x(y \text{Tr}_\ell^t(b) + b \text{Tr}_\ell^t(y)) + x(b \text{Tr}_\ell^t(b) + \alpha b^2)),$$

which is an affine function if and only if  $p(y) = y \text{Tr}_\ell^t(b) + b \text{Tr}_\ell^t(y)$  is constant for all  $y \in \mathbb{F}_{2^t}$ . If that is so, since  $p(0) = 0$ , then  $p(y) = y \text{Tr}_\ell^t(b) + b \text{Tr}_\ell^t(y) = 0$ , for all  $y$ , in particular, for  $y = 1$ , we get  $b + \text{Tr}_\ell^t(b) = 0$ , that is,

$$y \text{Tr}_\ell^t(b) + b \text{Tr}_\ell^t(y) = 0 \implies y + \text{Tr}_\ell^t(y) = 0 \implies y \in \mathbb{F}_{2^t}.$$

Thus  $p(y)$  is not a constant function for all  $y \in \mathbb{F}_{2^t}$ . Therefore  $g$  does not possess any affine derivative.  $\blacksquare$

**Remark 2.2:** If  $k = 1$ , then  $t = \ell$  and for any  $(a, b) \in \mathbb{F}_{2^t} \times \mathbb{F}_{2^t}$ ,

$$D_{(a,b)}g(x, y) = \text{Tr}_1^t((1 + \alpha)(xb^2 + a(y + b)^2)),$$

which is an affine function. Therefore, if  $k = 1$ , then a function  $g$  of the form as in Equation (17) has affine derivatives. Thus, if  $k = 1$ , then  $f_i$  and  $g$  are affine inequivalent, where  $f_i$  and  $g$  are defined as in (11) and (17), respectively.

**Theorem 2.4.** Suppose  $n = 2t$  and  $g$  be defined as in Equation (17). The number of distinct two-dimensional subspaces corresponding to constant second derivatives of  $g$  is

$$\frac{2^{-3\ell-1}(2^{2(2\ell+t)} + 2^{4\ell+t+1} - 2^{5\ell+t} - 5 \cdot 2^{3\ell+2t} + 2^{5\ell+2t} - 2^{2(\ell+t)} + 2^{3\ell+1} + 2^{4t})}{3}$$

**Proof:** Let  $V = \langle (a, b), (c, d) \rangle$  be any two-dimensional subspace of  $\mathbb{F}_{2^t} \times \mathbb{F}_{2^t}$ .

$$\begin{aligned} D_V g(x, y) &= \text{Tr}_1^t((ad + bc)\text{Tr}_\ell^t(y) + (a\text{Tr}_\ell^t(d) + c\text{Tr}_\ell^t(b))y + (ad\text{Tr}_\ell^t(d) + cb\text{Tr}_\ell^t(b)) \\ &\quad + (b\text{Tr}_\ell^t(d) + d\text{Tr}_\ell^t(b))x + (a + c)(b\text{Tr}_\ell^t(d) + d\text{Tr}_\ell^t(b)) + \alpha(ad^2 + cb^2)). \end{aligned}$$

*Case 1:* Let  $b = 0$  and  $d = 0$ . Then  $D_V g(x, y) = 0$ , for all  $(x, y) \in \mathbb{F}_{2^t} \times \mathbb{F}_{2^t}$ . Thus, with respect to any 2-dimensional subspace of  $\mathbb{F}_{2^t} \times \{0\}$ , the second order derivative of  $g$  is 0. The number of such two-dimensional subspaces is  $(2^t - 1)(2^{t-1} - 1)/3$ .

*Case 2:* Let  $b = 0$  and  $d \neq 0$ . Then

$$D_V g(x, y) = \text{Tr}_1^t(ad\text{Tr}_\ell^t(y) + a\text{Tr}_\ell^t(d)y + ad\text{Tr}_\ell^t(d) + \alpha ad^2). \quad (18)$$

Since

$$\begin{aligned} \text{Tr}_1^t(ad\text{Tr}_\ell^t(y)) &= \text{Tr}_1^t(ad(y + y^{2^\ell} + y^{2^{2\ell}} + \dots + y^{2^{(k-1)\ell}})) \\ &= \text{Tr}_1^t(y(ad + (ad)^{2^{(k-1)\ell}} + (ad)^{2^{(k-2)\ell}} + \dots + (ad)^{2^\ell})) \\ &= \text{Tr}_1^t(y\text{Tr}_\ell^t(ad)). \end{aligned}$$

From (18), we have

$$D_V g(x, y) = \text{Tr}_1^t((\text{Tr}_\ell^t(ad) + a\text{Tr}_\ell^t(d))y + ad\text{Tr}_\ell^t(d) + \alpha ad^2),$$

which is constant if and only if

$$\text{Tr}_\ell^t(ad) + a\text{Tr}_\ell^t(d) = 0. \quad (19)$$

*Subcase (i):* Let  $a \in \mathbb{F}_{2^t}$ . Then Equation (19) is satisfied for all  $d \in \mathbb{F}_{2^t}$ . Therefore,  $d$  can be chosen in  $2^t - 1$  ways and  $a$  in  $2^\ell - 1$  ways. Thus, the number of two-dimensional subspaces on which the second derivatives of  $g$  are constants is equal to  $(2^\ell - 1)2^{t-1}(2^t - 1)$ .

*Subcase (ii):* Let  $a \in \mathbb{F}_{2^t} \setminus \mathbb{F}_{2^\ell}$ . Then  $\text{Tr}_\ell^t(ad) + a\text{Tr}_\ell^t(d) = 0$  if and only if  $\text{Tr}_\ell^t(d) = 0$  and  $\text{Tr}_\ell^t(ad) = 0$ . Since both are  $(k-1)$ -dimensional  $\mathbb{F}_{2^\ell}$ -subspaces of  $\mathbb{F}_{2^t}$ ,  $d$  can be chosen in  $2^{t-2\ell}$  ways and  $a$  in  $2^t - 2^\ell$  ways. Thus, the number of such two-dimensional subspaces is  $(2^t - 2^\ell)2^{t-1}(2^{t-2\ell} - 1)$ .

*Case 3* Let  $b \neq 0$  and  $d \neq 0$  and  $b = d$ . Then the subspace generated by  $\{(a, b), (c, d)\}$  is equal to the subspace generated by  $\{(a + c, b + d), (c, d)\} = \{(a + c, 0), (c, d)\}$ , which is the same as in the previous case.

*Case 4:* Let  $b \neq 0$  and  $d \neq 0$  and  $b \neq d$ . Then  $D_V g(x, y)$  is constant if and only if

$$b\text{Tr}_\ell^t(d) + d\text{Tr}_\ell^t(b) = 0, \quad \text{for all } x \in \mathbb{F}_{2^t} \quad (20)$$

and we get the implications

$$\begin{aligned} \text{Tr}_1^t((ad + bc)\text{Tr}_\ell^t(y) + (a\text{Tr}_\ell^t(d) + c\text{Tr}_\ell^t(b))y) &= 0, \\ \text{Tr}_1^t((\text{Tr}_\ell^t(ad + bc) + (a\text{Tr}_\ell^t(d) + c\text{Tr}_\ell^t(b)))y) &= 0, \quad \text{for all } y \in \mathbb{F}_{2^t}, \\ \text{Tr}_\ell^t(ad + bc) &= a\text{Tr}_\ell^t(d) + c\text{Tr}_\ell^t(b). \end{aligned} \quad (21)$$

*Subcase (i):* Let  $\text{Tr}_\ell^t(b) = 0$  and  $\text{Tr}_\ell^t(d) = 0$ . The dimension of  $\ker(\text{Tr}_\ell^t)$  is  $t - \ell$ , where  $\ker(\text{Tr}_\ell^t) = \{x \in \mathbb{F}_{2^t} : \text{Tr}_\ell^t(x) = 0\}$ . Thus,  $d$  can be chosen in  $2^{t-\ell} - 1$  ways and  $b$  in  $2^{t-\ell} - 2$  ways. From Equation (21), we get  $\text{Tr}_\ell^t(ad + bc) = 0$ , so  $\text{Tr}_\ell^t(ad) = \text{Tr}_\ell^t(cb) = \lambda \in \mathbb{F}_{2^\ell}$ . For fixed  $b$  and  $d$  and for each  $\lambda \in \mathbb{F}_{2^\ell}$ ,  $a$  and  $c$  both can be chosen in  $2^{t-\ell}$  ways. Thus, the number of such distinct two-dimensional subspaces is  $2^{2t-\ell}(2^{t-\ell} - 1)(2^{t-\ell-1} - 1)/3$ .

**Table 1.** The number of two-dimensional subspaces on which the second derivative of the cubic Maiorana–McFarland bent functions  $f_j$  and  $g$  are constants.

	$n = 6$ $e = 1; \ell = 3$	$n = 10$ $e = 1; \ell = 5$	$n = 12$ $e = 2; \ell = 2$	$n = 18$ $e = 1; \ell = 9$	$e = 3; \ell = 3$
$n_1(e)$	35	651	12075	174251	3052203
$n_2(\ell)$	651	174251	53675	11453115051	25287339
$n = 20$ $e = 2; \ell = 2$		$e = 1; \ell = 3$	$n = 30$ $e = 3; \ell = 5$		$e = 5; \ell = 15$
3142315	879630115		12526594731		188614879915
2831415467	3775311936432811		6052134955691		192153583564270240

*Subcase (ii):* Let  $\text{Tr}_\ell^t(b) = 0$  but  $\text{Tr}_\ell^t(d) \neq 0$  or  $\text{Tr}_\ell^t(b) \neq 0$  but  $\text{Tr}_\ell^t(d) = 0$ . Then, from Equation (20),  $b = 0$  or  $d = 0$  respectively, which is impossible.

*Subcase (iii):* Let  $\text{Tr}_\ell^t(b) \neq 0$  and  $\text{Tr}_\ell^t(d) \neq 0$ . From Equation (20), we get

$$d = \frac{\text{Tr}_\ell^t(d)}{\text{Tr}_\ell^t(b)} b, \text{ that is, } d = \beta b, \text{ where } \beta = \frac{\text{Tr}_\ell^t(d)}{\text{Tr}_\ell^t(b)} \in \mathbb{F}_{2^\ell}^* \text{ and } \beta \neq 1.$$

For each  $b \in \mathbb{F}_{2^\ell}^*$ ,  $d$  can be chosen in  $2^\ell - 2$  ways. From Equation (21), we get

$$\text{Tr}_\ell^t(b(a\beta + c)) = (a\beta + c)\text{Tr}_\ell^t(b). \tag{22}$$

Equation (22) has a solution if and only if  $a\beta + c \in \mathbb{F}_{2^\ell}$ , so,  $c = a\beta + \beta_1$ , where  $\beta_1 \in \mathbb{F}_{2^\ell}$ . Then for any fixed  $a$ ,  $c$  can be chosen in  $2^\ell$  ways. Therefore the number of such two-dimensional distinct subspaces is  $2^{t+\ell}(2^t - 1)(2^{\ell-1} - 1)/3$ . Adding all the cases we get our count. ■

In what follows we demonstrate affine inequivalence among the cubic bent functions constructed above. To do this we use Theorem 1.3 proved in [11]. However, it is to be remembered that the use of the properties of higher-order derivatives to decide affine inequivalence between bent function was introduced by Dillon [9] way back in the seventies.

**Remark 2.3:** Let  $n = 2t$  be a fixed positive integer. In Theorem 2.4, we proved that the number of two-dimensional subspaces with respect to which the second-order derivatives of  $g$  are constants depends on  $\ell$ . Thus, for any fixed  $n$ , for different choices of  $\ell$  the number of distinct two-dimensional subspaces to constant second derivatives of the corresponding functions are different. Therefore, for any fixed  $n$ , for different choices of  $\ell$  the corresponding cubic Maiorana–McFarland bent functions are affine inequivalent.

**Example 2.1:** Let  $n = 30$ . Then  $t = 15$  and possible values of  $\ell$  are 3, 5, and 15. If  $\ell = 15$ , then  $k = 1$  and, the bent function corresponding to  $\ell = 15$  has an affine derivative. Also from Table 1, we get the cubic bent functions corresponding to  $\ell = 3$ ,  $\ell = 5$  and  $\ell = 15$  are mutually affine inequivalent.

Let  $n = 2t$ , and  $n_1(e)$  and  $n_2(\ell)$  be the number of two-dimensional subspaces of  $\mathbb{F}_{2^t} \times \mathbb{F}_{2^t}$  on which the second derivatives of  $f_i$  and  $g$  (defined as in Equations (11) and (17), respectively) are constants. Then

$$n_1(e) = \frac{(2^t - 1)(2^{t+e-1}(2^e + 1) - (2^t + 1))}{3} \tag{23}$$

and

$$n_2(\ell) = \frac{(2^t - 1)(2^{t+\ell}(2^{\ell-1} - 1) + 2^{t-1} - 1) + 2^{2t-\ell}(2^{t-\ell} - 1)(2^{t-\ell-1} - 1)}{3} + 2^{t-1}((2^\ell - 1)(2^t - 1) + (2^t - 2^\ell)(2^{t-2\ell} - 1)). \quad (24)$$

**Lemma 2.2.** *If  $\ell \geq e$ , then  $n_2(\ell) > n_1(e)$ , where  $n_1(e)$  and  $n_2(\ell)$  are defined as in Equations (23) and (24), respectively.*

**Proof:** We first compute the difference

$$\begin{aligned} n_2(\ell) - n_1(e) &= (2^\ell - 1)2^{t-1}(2^t - 1) + (2^t - 2^\ell)2^{t-1}(2^{t-2\ell} - 1) \\ &\quad - (2^t - 1)2^{t-1}(2^e - 1) + \frac{2^{2t-\ell}(2^{t-\ell} - 1)(2^{t-\ell-1} - 1)}{3} \\ &\quad + \frac{2^{t+\ell}(2^t - 1)(2^{\ell-1} - 1)}{3} - \frac{2^{t+e}(2^t - 1)(2^{e-1} - 1)}{3}. \end{aligned}$$

If  $\ell = e$ , then  $\ell < t$  and  $n_2(\ell) - n_1(e) > 0$ . Again if  $\ell > e$  then  $n_2(\ell) - n_1(e) > 0$  since

$$(2^\ell - 1)2^{t-1}(2^t - 1) - (2^t - 1)2^{t-1}(2^e - 1) > 0$$

and

$$\frac{2^{t+\ell}(2^t - 1)(2^{\ell-1} - 1)}{3} - \frac{2^{t+e}(2^t - 1)(2^{e-1} - 1)}{3} > 0. \quad \blacksquare$$

**Corollary 2.1.** *If  $\ell \geq e$ , then  $f_\ell$  and  $g$  are affine inequivalent, where  $f_\ell$  and  $g$  are defined as in (11) and (17), respectively.*

We compare  $n_1(e)$  and  $n_2(\ell)$  in Table 1, for different values of  $n$ .

### 3. Conclusion

In this article, we prove that cubic Maiorana–McFarland bent functions which are constructed by using some known types of permutation polynomials (see [1, 13]) have no affine derivative. Consequently, we have obtained many affine inequivalent classes of bent function within the functions under consideration.

### Acknowledgments

The authors thanks the anonymous referees and the editor for useful comments that has improved the technical and editorial quality of the article.

### Disclosure statement

No potential conflict of interest was reported by the authors.

### Funding

Bimal Mandal thanks IIT Roorkee for the research grant.



## References

- [1] A. Blokhuis, R.S. Coulter, M. Henderson and C.M. O'Keefe, *Permutations amongst the Dembowski-Ostrom polynomials*, in *1999 Finite Fields and Applications*, D. Jungnickel and H. Niederreiter, eds., Springer, Berlin, 2001, pp. 37–42.
- [2] P. Camion, C. Carlet, P. Charpin, and N. Sendrier, *On correlation-immune functions*, in *Advances in Cryptology - CRYPTO '91*, CRYPTO 1991, LNCS 576, J. Feigenbaum, ed., Springer, Berlin, 1992, pp. 86–100.
- [3] A. Canteaut and P. Charpin, *Decomposing bent functions*, IEEE Trans. Inform. Theory 49(8) (2003), pp. 2004–2019.
- [4] C. Carlet, *Two new classes of bent functions*, in *Advances in Cryptology - EUROCRYPT '93*, EUROCRYPT 1993, LNCS 765, T. Helleseth, ed., Springer, Berlin, 1994, pp. 77–101.
- [5] C. Carlet, *Recursive lower bounds on the nonlinearity profile of Boolean functions and their applications*, IEEE Trans. Inform. Theory 54(3) (2008), pp. 1262–1272.
- [6] P. Charpin and S. Sarkar, *Polynomials with Linear Structure and Maiorana-McFarland Construction*, IEEE Trans. Inform. Theory 57(6) (2011), pp. 3796–3804.
- [7] R.S. Coulter, *On the evaluation of a class of Weil sums in characteristic 2*, New Zealand J. Math 28 (1999), pp. 171–184.
- [8] T.W. Cusick and P. Stănică, *Cryptographic Boolean Functions and Applications*, Academic Press, San Diego, CA, 2009.
- [9] J.F. Dillon, *Elementary Hadamard Difference Sets*, Proceedings of 6th S. E. Conference on Combinatorics, Graph Theory, and Computing, Utility Mathematics, Winnipeg, 1975, pp. 237–249.
- [10] S. Gangopadhyay, S. Sarkar, and R. Telang, *On the lower bounds of the second order nonlinearities of some Boolean functions*, Inform. Sci. 180 (2010), pp. 266–273.
- [11] S. Gangopadhyay, *Affine inequivalence of cubic Maiorana-McFarland type bent functions*, Discret. Appl. Math 161 (2013), pp. 1141–1146.
- [12] X.-D. Hou, *Cubic bent functions*, Discret. Math 189 (1998), pp. 149–161.
- [13] Y. Lalgè-Chapuy, *A Note on a Class of Quadratic Permutations Over  $\mathbb{F}_2^n$* , in *Applied Algebra, Algebraic Algorithms and Error-Correcting Codes. AAEECC 2007*, LNCS 4851, S. Boztaş and H.F. Lu, eds., Springer, Berlin, pp. 130–137.
- [14] E.J. MacWilliams and N.J.A. Sloane, *The Theory of Error-Correcting Codes*, North-Holland, Amsterdam, 1977.
- [15] O.S. Rothaus, *On bent functions*, J. Combin. Theory - Ser. A 20 (1976), pp. 300–305.
- [16] T. Stengenthaler, *Correlation-immunity of nonlinear combining functions for cryptographic applications*, IEEE Trans. Inf. Theory 30(5) (1984), pp. 776–780.