



Calhoun: The NPS Institutional Archive

DSpace Repository

Reports and Technical Reports

All Technical Reports Collection

1978-06

Simulation of Nonhomogeneous Poisson Processes by Thinning

Lewis, P.A.W.; Shedler, G.S.

Monterey, California. Naval Postgraduate School

https://hdl.handle.net/10945/63247

This publication is a work of the U.S. Government as defined in Title 17, United States Code, Section 101. Copyright protection is not available for this work in the United States.

Downloaded from NPS Archive: Calhoun



Calhoun is the Naval Postgraduate School's public access digital repository for research materials and institutional publications created by the NPS community. Calhoun is named for Professor of Mathematics Guy K. Calhoun, NPS's first appointed -- and published -- scholarly author.

> Dudley Knox Library / Naval Postgraduate School 411 Dyer Road / 1 University Circle Monterey, California USA 93943

http://www.nps.edu/library



NPS-55-78-014 NAVAL POSTGRADUATE SCHOOL Monterey, California



OCT 18

SIMULATION OF NONHOMOGENEOUS POISSON PROCESSES BY THINNING

by

P. A. W. Lewis and G. S. Shedler

JUNE 1978

Approved for public release; distribution unlimited.

PUC FILE COPY AD AO 59904

NAVAL POSTGRADUATE SCHOOL MONTEREY, CALIFORNIA

Rear Admiral Tyler F. Dedman Superintendent

Jack R. Borsting Provost

Reproduction of all or part of this report is authorized.

This report was prepared by:

Ters a.W. Le

Peter A. W. Lewis, Professor Department of Operations Research

by PAWL.

Gerald S. Shedler IBM Research Laboratory

Reviewed by:

Released by:

. Idlen William M. Tolles

Acting Dean of Research

Michael G. Sovereign, Chairman

Department of Operations Research

1. GOVT ACCESSION NO. 1. RECIPIENT'S CATALOG NUMBER NPESS-78-14 2. GOVT ACCESSION NO. 1. RECIPIENT'S CATALOG NUMBER NPESS-78-14 3. TTHE of RESORT & PERIOD COVER 5. THE OF REPORT A PERIOD COVER 5. THE OF REPORT AND THE OFFICE STATEMENT NUMBER 5. THE OF REPORT AND THE OFFICE STATEMENT NUMBER 5. THE OF REPORT AND THE OFFICE STATEMENT NUMBER 5. THE OF REPORT AND THE OFFICE STATEMENT NUMBER 5. THE OF REPORT AND THE OFFICE STATEMENT NUMBER 5. THE OF REPORT AND THE OFFICE STATEMENT OF THE OFFICE STATEMENT OFFICE STATEMENT OF THE OFFICE STATEMENT OF THE OFFICE STATEMENT OFFICE STATEMENT OF THE OFFICE STATEMENT OF THE OFFICE STATEMENT OFFICE STATEMENT OF THE OFFICE ST	1. PERAT-NUMBER	READ INSTRUCTIONS BEFORE COMPLETING FORM	
 TITLE (and Submit) TITLE (and Submit) Simulation of Nonhomogeneous Poisson Processes Multipace Peter A. W. Jewis and Grow Shedler Peter A. W. Jewis And And Access Office of Naval Research Arlington, VA 22217 Nonotrou No Or Aut And Access Office of Naval Research M. Junger B. Jewis And Access Office of Naval Research W. M. K. D. J. H. S. Jewis And Access Office of Path Act And Access (distribution unlimited) W. M. K. D. J. H. S. Jewis Access (distribution unlimited) W. M. K. D. Jewis Access (distribution unlimited) W. M. K. D. Jewis Access (distribution unlimited) W. K. W. W. M. K. Accesses, two-dimensional nonhomogeneous Poisson processes, one-dimensional nonhomogeneous Poisson processes, thinning, search and detection, acceptance-rejection, order statistics. A simple and relatively efficient method for simulating one-dimensional an two-dimensional nonhomogeneous Poisson processes is presented. The method is applicable for any rate function and is based on controlled deletion of points in a Poisson processe whose rate function, for ordering of points, and fo generation of Poisson variates. M. K. W. W. M. K. M. Statisti and the subscription without of points, and fo generation of Poisson variates. 	NPS55-78-14	0. 3. RECIPIENT'S CATALOG NUMBER	
 A TITLE Card Sketting Simulation of Nonhomogeneous Poisson Processes ANTHORNAL CARD ALL AND ADDRESS Peter A. W. Lewis and OFFEY: Shedler CONTRACT ON GRANT NUMBER() Peter A. W. Lewis and OFFEY: Shedler CONTRACT ON GRANT NUMBER() Peter A. W. Lewis and OFFEY: Shedler CONTRACT ON GRANT NUMBER() Peter A. W. Lewis and OFFEY: Shedler CONTRACT ON GRANT NUMBER() Peter A. W. Lewis and OFFEY: Shedler CONTRACT ON GRANT NUMBER() Peter A. W. Lewis and OFFEY: Shedler CONTRACT ON GRANT NUMBER() Peter A. W. Lewis and OFFEY: Shedler CONTRACT ON GRANT NUMBER() Peter A. W. Lewis and OFFEY: Shedler CONTRACT ON GRANT NUMBER() CONTRACT ON GRANT NUMBER() CONTRACT ON GRANT NUMBER() CONTRACT ON GRANT NUMBER() CONTRACT ON AND ADDRESS Office of Naval Research Arlington, VA 22217 NON170K-G ADENCY MARE ADDRESS() different from Controlling Office M. MUMERT OFFICE NAME ADDRESS() different from Controlling Office M. MUMERT OFFICE NAME ADDRESS() different from Controlling Office M. MUMERT OFFICE NAME ADDRESS() different from Repert() M. STATEMENT (of the abstract entered in Block 20, if different from Repert() T. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Repert() SUPPLEMENTARY NOTES SUPPLEMENTARY NOTES SUPPLEMENTARY NOTES A SIGN Continue on reverse side if necessary and Identify by Moch number() A SIGN Continue on reverse side if necessary and Identify by Moch number() A SIGN Continue on reverse side if necessary and Identify by Moch number() Approved for public release; the different from Repert() SUPPLEMENTARY NOTES SUPPLEMENTARY NOTES SUPPLEMENTARY NOTES A SIGN Processes, the different from Repert() A SIGN Processes, simulati			
 PERFORMING ORGANIZATION ROME FOR Sev Shedler PERFORMING ORGANIZATION ROME FOR Sev Shedler PERFORMING ORGANIZATION ROME FOR Sev Shedler CONTRACT ON GRANT NUMBER/OF Sev Shedler CONTRACT ON GRANT NUMBER/OF Sev Shedler PERFORMING ORGANIZATION ROME FOR Sev Shedler PERFORMING ORGANIZATION ROME FOR Several Se	4. TITLE (and Subtitle)	5. TYPE OF REPORT & PERIOD COVERE	
ALTIMERA Peter A. W Lewis and GFB: Shedler Peter A. W Lewis and A Dottess Office of Naval Research Arlington, VA 22217 Wontforwer Addition A Dottess Office of Naval Research Arlington, VA 22217 BECHATION OFFICE AND A DOTTES WONGER OF ANY LABORESS (I dilerent from Controlling Office) BECHATON A DOTTAGE ADDRESS Office of Naval Research Approved for public release; distribution unlimited. D FFR: D P A D A DOTTAGE ADDRESS (I dilerent from Report) Unclassified BECHATON FOR FAREWARD A DOTTAGE ADDRESS Office of public release; distribution unlimited. D FFR: D P A D A DOTTAGE ADDRESS (I dilerent from Report) D TO DISTRIBUTION STATEMENT (of the abstract entreed in Block 20, II dilerent from Report) T. DISTRIBUTION STATEMENT (of the abstract entreed in Block 20, II dilerent from Report) Poisson processes, nonhomogeneous Poisson processes, one-dimensional non-homogeneous Poisson processes, thunding, search and detection, acceptance-rejection, order statistics. Astract (Continue on reverse ald II necessary and Identify by Block number) A simple and relatively efficient method for simulating one-dimensional an two-dimensional nonhomogeneous Poisson processes is presented. The method is applicable for any rate function and is based on control delection, acceptance-rejection of the rate function dis based on control delection of points in	by Thinning	Giechnical PEPIS	
AUTION OF ACT ON GRANT NUMBERS Peter A. W./Lewis mid-Grads-Sr-Shedler (SEPA) (Signature School Monterey, Ca. 93940 PENDAMING ORGANIZATION THAT AND ABORESS Office of Naval Research Arlington, VA 22217 It controlling Office NAME AND ADDRESS Office of Naval Research Arlington, VA 22217 It MONITON''G AGENCY HAME A ADDRESS Office of Naval Research Arlington, VA 22217 It MONITON''G AGENCY HAME A ADDRESS Office of Naval Research Arlington, VA 22217 It MONITON''G AGENCY HAME A ADDRESS Office of Naval Research Arlington, VA 22217 It MONITON''G AGENCY HAME A ADDRESS Office of Naval Research Arlington, VA 22217 It MONITON''G AGENCY HAME A ADDRESS Office of Naval Research Arlington, VA 22217 It MONITON''G AGENCY HAME A ADDRESS Office of naval Research Arlington, VA 22217 It MONITON''G AGENCY HAME A ADDRESS Office of naval Research Approved for public release; distribution unlimited. It SUPPLEMENTARY NOTES Asimple and relatively efficient method for simulating one-dimensional non-homogeneous Poisson processes, thoread on detection, acceptance-rejection, order statistics. A simple and relatively efficient method for simulating one-dimensional an two-dimensional nonhomogeneous Poisson processes is presented. The method is applicable for any rate function and is based on controlled deletion of points in a Poisson process whose rate function dominates the given rate function. In its simplest implementation, the method obviates the need for numerical integration of the rate function, for ordering of points, and for generation of Poinson variates. DD 100071 1473 connor 'Novel is obsolver'E	and a second	6. PERFORMING ORG. REPORT NUMBER	
Peter A. W. Jewis and G. S. Sheller (PEPONING ORGANIZATION FINE AND ABORESS Naval Postgraduate School Monterey, Ca. 93940 I. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Arlington, VA 22217 I. NONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Arlington, VA 22217 I. NONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Arlington, VA 22217 I. NONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Arlington, VA 22217 I. NONTOR-WG ADDRESS Office of Naval Research Approved for public release; distribution unlimited. I. SUPPLEMENTARY NOTES I. SUP	7. AUTHORICAL	S. CONTRACT OR GRANT NUMBER(.)	
9 FERFORMING ORDANIZATION THE AND ADDRESS 0 AREA N FORK UNIT NUMBERS Naval Postgraduate School 61153N, RR014-05-01 N0001478WR80035 In: CONTROLLING OFFICE NAME AND ADDRESS 61153N, RR014-05-01 N0001478WR80035 In: CONTROLLING OFFICE NAME AND ADDRESS 61153N, RR014-05-01 N0001478WR80035 In: CONTROLLING OFFICE NAME AND ADDRESS 61153N, RR014-05-01 N0001478WR80035 In: CONTROLLING OFFICE NAME AND ADDRESS 61153N, RR014-05-01 N0001478WR80035 II: CONTROLLING OFFICE NAME & AND ADDRESS III: CONTROLLING OFFICE NAME & ADDRESS(II dillerent from Controlling Office) III: SECURITY CLASS, (of this report) IV: DONTOR ''G AGENCY NAME & ADDRESS(II dillerent from Controlling Office) III: SECURITY CLASS, (of this report) IV: DESTRIBUTION STATEMENT (of the abalised entered in Black 20, II dillerent from Report) Unclassified IV: DISTRIBUTION STATEMENT (of the abalised entered in Black 20, II dillerent from Report) III: SUPPLEMENTARY NOTES IV: DISTRIBUTION STATEMENT (of the abalised entered in Black 20, II dillerent from Report) III: SUPPLEMENTARY NOTES IV: SUPPLEMENTARY NOTES III: SUPPLEMENTARY NOTES IV: SUPPLEMENTARY NOTES III: SUPPLEMENTARY NOTES IV: SUPPLEMENTARY NOTES IIII: SUPPLEMENTARY NOTES IV: SUPPLEMENTARY NO	Peter A. W. Lewis and Grass Shedler		
 FERFORMING CRANNEL AND ADDRESS Monterey, Ca. 93940 CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Arlington, VA 22217 CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Arlington, VA 22217 CONTROLLING OFFICE NAME A ADDRESS Office of Naval Research Arlington, VA 22217 CONTROLLING OFFICE NAME A ADDRESS(II different from Controlling Office) F. R. D. 1405 J. D. SECURITY CLASS. (<i>it into report</i>) U. R. R. D. 1405 J. D. SECURITY CLASS. (<i>it into report</i>) U. C. LASS. (<i>it into report</i>) J. D. SECURITY CLASS. (<i>it into report</i>) J. D. SECURITY FOR SECURITY (<i>it into an investe alds II necessary and leavely by Nock number</i>) POISON processes, nonhomogeneous Poisson processes, thronding, search and detection, acceptance-rejection, order statistics. J. ASSTRACT (Continue on reverse alds II necessary and Identify by Nock number) A SIMPLE MENTARY NOTES A SIMPLE and relatively efficient method for simulating one-dimensional and two-dimensional nonhomogeneous Poisson processes is presented. The method is applicable for any rate function and is based on controlled deletion of points in a Poisson process whose rate function dominates the given rate function. In the simplest implementation, the method obviates the need for numerical integration of the rate function, for ordering of points, and for generation of Points of the rate function, for ordering of points, and for generation of Points and Function. 	Gerald Silshed H		
Naval Postgraduate School Monterey, Ca. 93940 61153N, RR014-05-01 N 0001478WR80035 II. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Arlington, VA 22217 III. Security CLASS. (ci this report) Unclassified II. MONITOR & AGENCY MARE & ADDRESS Office of Naval Research Arlington, VA 22217 III. SECURITY CLASS. (ci this report) Unclassified III. DECLASS. (CI this report) III. SECURITY CLASS. (ci this report) III. DECLASS. (CATION/DOWNGRADIN) Approved for public release; distribution unlimited. III. DISTRIBUTION STATEMENT (ci the ebetrect entered in Block 20, If different from Report) III. SUPPLEMENTARY NOTES	9 PERFORMING ORGANIZATION NAME AND ADDRESS		
 N 0001478WR80035 N 0001478WR80035 N 0001478WR80035 Office of Naval Research Arlington, VA 22217 WONTON '' GAGENEY MARE & ADDRESS(II different from Controlling Office) SECURITY CLASS. (of this report) URE SECURITY CLASS. (of this report) Unclassified MRAL 485 OF FACES MARE A ADDRESS(II different from Controlling Office) SECURITY CLASS. (of this report) Unclassified DRARL 485 OF FACES Approved for public release; distribution unlimited. DF ATTEMENT (of the abstract entered in Block 20, If different from Report) IS DEFINED TO POSES IS DEFINISUTION STATEMENT (of the abstract entered in Block 20, If different from Report) IS DEFINISUTION STATEMENT (of the abstract entered in Block 20, If different from Report) IS DEFINISUTION STATEMENT (of the abstract entered in Block 20, If different from Report) IS DEFINISUTION STATEMENT (of the abstract entered in Block 20, If different from Report) IS DEFINISUTION STATEMENT (of the abstract entered in Block 20, If different from Report) IS DEFINISUTION OF TAKES IS ASTRACT (Continue on reverse side if inscenses and identify by block number) A SIGNAL TO FORME ADDRESS (INCOMENT ADDRESS) ABSTRACT (Continue on reverse side if inscenses and identify by block numbe	Monterey, Ca. 93940	6]153N, RR014-05-0]	
 11. CONTROLLING OFFICE NAME AND ADDRESS OFFICE of Naval Research Arlington, VA 22217 12. MONITOR CARENCY NAME & ADDRESS (II dillormi from Controlling Office) UNAVER OF FACE 27 UNCLassified 13. SECURITY CLASS. (of this report) UNCLassified 14. DECLASSIFICATION/DOWNGRADIN SCHEDULE 15. NEY WORDS (Continue on reverse side if necessary and identify by block number) Poisson processes, nonhomogeneous Poisson processes, one-dimensional non- homogeneous Poisson processes, two-dimensional nonhomogeneous Poisson processes, simulation of point processes, thinning, search and detection, acceptance-rejection, order statistics. 13. ABSTRACT (Continue on reverse side if necessary and identify by block number) A simple and relatively efficient method for simulating one-dimensional an two-dimensional nonhomogeneous Poisson processes is presented. The method is applicable for any rate function and is based on controlled deletion of points in a Poisson process whose rate function dis based on controlled deletion of points in a Poisson variates. DD 1/100/13. TATTA TEMENT of the rate function, for ordering of points, and fo generation of Poisson variates. 		N 0001478WR80035	
WILLE OF HAVAL RESERVENT Arlington, VA 22217 11. MONITOR OF AGENCY MARE & ADDRESS(II dillerent from Controlling Diller) 12. MONITOR OF AGENCY MARE & ADDRESS(II dillerent from Controlling Diller) 13. SECURITY CLASS. (of this report) 13. SECURITY CLASS. (of this report) 14. MONITOR OF AGENCY MARE & ADDRESS(II dillerent from Controlling Diller) 15. SECURITY CLASS. (of this report) 16. DITTRIBUTION STATEMENT (of the ebetract intered in Black 20, II dillerent from Report) 17. DISTRIBUTION STATEMENT (of the ebetract intered in Black 20, II dillerent from Report) 18. SUPPLEMENTARY NOTES 19. ASITY ACT (Continue on reverse side If necessary and identify by block number) Poisson processes, nonhomogeneous Poisson processes, one-dimensional non-homogeneous Poisson processes, two-dimensional nonhomogeneous Poisson processes, simulation of point processes, thinning, search and detection, acceptance-rejection, order statistics. 13. ASITYACT (Continue on reverse side II necessary and identify by block number) A simple and relatively efficient method for simulating one-dimensional an two-dimensional nonhomogeneous Poisson processes is presented. The method is applicable for any rate function and is based on controlled deletion of points in a Poisson process whose rate function dominates the given rate f	11. CONTROLLING OFFICE NAME AND ADDRESS	12 REPORT DATE	
 17. MONITOR''' G AGENCY NAME & ADDRESS(II dillorent from Controlling Office) 18. SECURITY CLASS. (of this report) Unclassified 19. DECLASSIFICATION/DOWNGRADIN 10. DECLASSIFICATION/DOWNGRADIN 10. DECLASSIFICATION/DOWNGRADIN 11. DISTRIBUTION STATEMENT (of the observed in Block 20, If dillorent from Report) 12. DISTRIBUTION STATEMENT (of the observed in Block 20, If dillorent from Report) 19. KEY WORDS (Continue on reverse of the II necessary and Identify by Noch number) Poisson processes, nonhomogeneous Poisson processes, one-dimensional nonhomogeneous Poisson processes, two-dimensional nonhomogeneous Poisson processes, thinning, search and detection, acceptance-rejection, order statistics. 20. ABSTRACT (Continue on reverse of the II necessary and Identify by Noch number) A simple and relatively efficient method for simulating one-dimensional an two-dimensional nonhomogeneous Poisson processes is presented. The method is applicable for any rate function and is based on controlled deletion of points in a Poisson process whose rate function dominates the given rate function. In its simplest implementation, the method obviates the need for numerical integration of the rate function, for ordering of points, and fo generation of Poisson variates. 	Arlington, VA 22217	11 JULINES / O	
 14. MONITOR^{WIG} AGENCY MAMER ADDRESS(II different from Controlling Office) 15. SECURITY CLASS. (of this report) Unclassified 16. DECLASSIFICATION/DOWNGRADIN 17. DISTRIBUTION STATEMENT (of the observed in Block 20, if different from Report) 17. DISTRIBUTION STATEMENT (of the observed on Block 20, if different from Report) 18. SUPPLEMENTARY NOTES 19. ABSTRACT (Continue on reverse aids if necessary and identify by block number) Poisson processes, nonhomogeneous Poisson processes, one-dimensional nonhomogeneous Poisson processes, two-dimensional nonhomogeneous Poisson processes, two-dimensional nonhomogeneous Poisson processes, two-dimensional nonhomogeneous Poisson processes is presented. The method is applicable for any rate function and is based on controlled deletion of points in a Poisson process whose rate function dominates the given rate function. In its simplest implementation, the method obviates the need for numerical integration of the rate function, for ordering of points, and for generation of Poisson variates. 20. ABSTRACT (Continue on reverse side in processes whose rate function dominates the given rate function. In its simplest implementation, the method obviates the need for numerical integration of the rate function, for ordering of points, and for generation of Points is obsolver. 		27	
 WIT K D 1 405 1 WR R J 405 AL DR R R R J 405 AL DR R R R J 405 AL DR R R R J 405 AL DISTRIBUTION STATEMENT (of the obstract entered in Black 20, if different from Report) DISTRIBUTION STATEMENT (of the obstract entered in Black 20, if different from Report) DISTRIBUTION STATEMENT (of the obstract entered in Black 20, if different from Report) SUPPLEMENTARY NOTES SUPPLEMENTARY NOTES SUPPLEMENTARY NOTES A SUPPLEMENTARY NOTES A SUPPLEMENTARY NOTES A Simple and relatively efficient method for simulating one-dimensional and two-dimensional nonhomogeneous Poisson processes is presented. The method is applicable for any rate function and is based on controlled deletion of points in a Poisson process whose rate function dominates the given rate function. In its simplest implementation, the method obviates the need for numerical integration of the rate function, for ordering of points, and for generation of Poisson variates. DD 1007 1 1473 EDITION OF 1 NOV 65 IS OBSOLETE 	14. MONITORING AGENCY NAME & ADDRESS(II dillerent from Controlling Office,) 15. SECURITY CLASS. (of this report)	
 16. DECLASSIFICATION/DOWNGRADIN 17. DISTRIBUTION STATEMENT (of the observed entered in Block 20, if different from Report) 18. SUPPLEMENTARY NOTES 19. SUPPLEMENTARY NOTES 19. SUPPLEMENTARY NOTES 19. ABSTRACT (Continue on reverse side if necessary and identify by block number) Poinson processes, nonhomogeneous Poinson processes, one-dimensional non-homogeneous Poisson processes, thinning, search and detection, acceptance-rejection, order statistics. 20. ABSTRACT (Continue on reverse side if necessary and identify by block number) A simple and relatively efficient method for simulating one-dimensional an two-dimensional nonhomogeneous Poisson processes is presented. The method is applicable for any rate function and is based on controlled deletion of points in a Poisson process rate function dominates the given rate function. In its simplest implementation, the method obviates the need for numerical integration of the rate function, for ordering of points, and for generation of Poisson variates. 20. Juny 71 1473 EDITION of 1 NOV 65 IS OBSOLETE 	16/ F. R. D 1 7 05	000102011160	
 16. Dity biournow statement (of the Report) Approved for public release; distribution unlimited. 17. DISTRIBUTION STATEMENT (of the ebetract entered in Block 20, if different from Report) 17. DISTRIBUTION STATEMENT (of the ebetract entered in Block 20, if different from Report) 18. SUPPLEMENTARY NOTES 19. SUPPLEMENTARY NOTES 19. KEY WORDS (Continue on reverse elds if necessary and identify by block number) Poisson processes, nonhomogeneous Poisson processes, one-dimensional non-homogeneous Poisson processes, two-dimensional nonhomogeneous Poisson processes, thunling, search and detection, acceptance-rejection, order statistics. 20. Asstract (Continue on reverse elds if necessary and identify by block number) A simple and relatively efficient method for simulating one-dimensional an two-dimensional nonhomogeneous Poisson processes is presented. The method is applicable for any rate function and is based on controlled deletion of points in a Poisson process whose rate function dominates the given rate function. In its simplest implementation, for ordering of points, and fo generation of Poisson variates. DD 1407.73 1473 EDITION OF 1 NOV 68 IS OBSOLETE 	MRRØ140501	15e, DECLASSIFICATION/DOWNGRADING	
Approved for public release; distribution unlimited.	16. DISTRIBUTION STATEMENT (of this Report)	. 1 1	
 W H3 pi 17. DISTRIBUTION STATEMENT (of the observed in Block 20, 11 different from Report) 18. SUPPLEMENTARY NOTES 19. KEY WORDS (Continue on reverse side II necessary and identify by block number) Poisson processes, nonhomogeneous Poisson processes, one-dimensional non- homogeneous Poisson processes, two-dimensional nonhomogeneous Poisson pro- cesses, simulation of point processes, thinning, search and detection, acceptance-rejection, order statistics. 28. ABSTRACT (Continue on reverse side II necessary and identify by block number) A simple and relatively efficient method for simulating one-dimensional an two-dimensional nonhomogeneous Poisson processes is presented. The method is applicable for any rate function and is based on controlled deletion of points in a Poisson process whose rate function dominates the given rate function. In its simplest implementation, the method obviates the need fo numerical integration of the rate function, for ordering of points, and fo generation of Poisson variates. DD 1100 73 1473 EDITION OF 1 NOV 64 IS OBSOLETE 	Approved for public release; distribution unlin	11tea.	
 17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) 18. SUPPLEMENTARY NOTES 19. KEY WORDS (Continue on reverse aids if necessary and identify by block number) Poisson processes, nonhomogeneous Poisson processes, one-dimensional non-homogeneous Poisson processes, two-dimensional nonhomogeneous Poisson processes, simulation of point processes, thinning, search and detection, acceptance-rejection, order statistics. 28. ABSTRACT (Continue on reverse aids if necessary and identify by block number) A simple and relatively efficient method for simulating one-dimensional an two-dimensional nonhomogeneous Poisson processes is presented. The method is applicable for any rate function and is based on controlled deletion of points in a Poisson process whose rate function dominates the given rate function. In its simplest implementation, the method obviates the need fo numerical integration of the rate function, for ordering of points, and fo generation of Poisson variates. 29. Norma 1473 EDITION OF INOV 68 IS OBSOLETE 	12, 4 3 p.		
 18. SUPPLEMENTANY NOTES 19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Poisson processes, nonhomogeneous Poisson processes, one-dimensional non- homogeneous Poisson processes, two-dimensional nonhomogeneous Poisson pro- cesses, simulation of point processes, thinning, search and detection, acceptance-rejection, order statistics. 28. ANSTRACT (Continue on reverse side if necessary and identify by block number) A simple and relatively efficient method for simulating one-dimensional an two-dimensional nonhomogeneous Poisson processes is presented. The method is applicable for any rate function and is based on controlled deletion of points in a Poisson process whose rate function dominates the given rate function. In its simplest implementation, the method obviates the need fo numerical integration of the rate function, for ordering of points, and fo generation of Poisson variates. DD 14AN 73 EDITION OF I NOV 68 IS OBSOLETE 	17. DISTRIBUTION STATEMENT (of the abetract entered in Block 20, if different	from Report)	
 18. SUPPLEMENTARY NOTES 19. KEY WORDS (Continue on reverse elde II necessary and identify by block number) Poisson processes, nonhomogeneous Poisson processes, one-dimensional non- homogeneous Poisson processes, two-dimensional nonhomogeneous Poisson pro- cesses, simulation of point processes, thinning, search and detection, acceptance-rejection, order statistics. 29. ABSTRACT (Continue on reverse elde II necessary and Identify by block number) A simple and relatively efficient method for simulating one-dimensional an two-dimensional nonhomogeneous Poisson processes is presented. The method is applicable for any rate function and is based on controlled deletion of points in a Poisson process whose rate function dominates the given rate function. In its simplest implementation, the method obviates the need fo numerical integration of the rate function, for ordering of points, and fo generation of Poisson variates. DD 1/AN 73 1473 EDITION OF 1 NOV 68 IS OBSOLETE 			
 18. SUPPLEMENTARY NOTES 19. KEY WORDS (Continue on reverse elde if necessery and identify by block number) Poisson processes, nonhomogeneous Poisson processes, one-dimensional non- homogeneous Poisson processes, two-dimensional nonhomogeneous Poisson pro- cesses, simulation of point processes, thinning, search and detection, acceptance-rejection, order statistics. 29. ABSTRACT (Continue on reverse elde if necessery and identify by block number) A simple and relatively efficient method for simulating one-dimensional an two-dimensional nonhomogeneous Poisson processes is presented. The method is applicable for any rate function and is based on controlled deletion of points in a Poisson process whose rate function dominates the given rate function. In its simplest implementation, the method obviates the need fo numerical integration of the rate function, for ordering of points, and fo generation of Poisson variates. DD 1/AN 73 1473 EDITION OF 1 NOV 60 15 OBSOLETE 			
 18. SUPPLEMENTARY NOTES 19. KEY WORDS (Continue on reverse elde II necessary and identify by block number) Poisson processes, nonhomogeneous Poisson processes, one-dimensional non- homogeneous Poisson processes, two-dimensional nonhomogeneous Poisson pro- cesses, simulation of point processes, thinning, search and detection, acceptance-rejection, order statistics. 28. ABSTRACT (Continue on reverse elde II necessary and identify by block number) A simple and relatively efficient method for simulating one-dimensional an two-dimensional nonhomogeneous Poisson processes is presented. The method is applicable for any rate function and is based on controlled deletion of points in a Poisson process whose rate function dominates the given rate function. In its simplest implementation, the method obviates the need fo numerical integration of the rate function, for ordering of points, and fo generation of Poisson variates. DD 1/AM 73 1473 EDITION OF 1 NOV 68 IS OBSOLETE 			
 18. KEY WORDS (Continue on reverse side if necessary and identify by block number) Poisson processes, nonhomogeneous Poisson processes, one-dimensional non- homogeneous Poisson processes, two-dimensional nonhomogeneous Poisson pro- cesses, simulation of point processes, thinning, search and detection, acceptance-rejection, order statistics. 28. ABSTRACT (Continue on reverse side if necessary and identify by block number) A simple and relatively efficient method for simulating one-dimensional an two-dimensional nonhomogeneous Poisson processes is presented. The method is applicable for any rate function and is based on controlled deletion of points in a Poisson process whose rate function dominates the given rate function. In its simplest implementation, the method obviates the need fo numerical integration of the rate function, for ordering of points, and fo generation of Poisson variates. DD tion 73 1473 EDITION OF 1 NOV 66 IS OBSOLETE 	18. SUPPLEMENTARY NOTES		
 19. KEY WORDS (Continue on reverse elde II necessary and identify by block number) Poisson processes, nonhomogeneous Poisson processes, one-dimensional non- homogeneous Poisson processes, two-dimensional nonhomogeneous Poisson pro- cesses, simulation of point processes, thinning, search and detection, acceptance-rejection, order statistics. 20. ABSTRACT (Continue on reverse elde II necessary and identify by block number) A simple and relatively efficient method for simulating one-dimensional an two-dimensional nonhomogeneous Poisson processes is presented. The method is applicable for any rate function and is based on controlled deletion of points in a Poisson process whose rate function dominates the given rate function. In its simplest implementation, the method obviates the need fo numerical integration of the rate function, for ordering of points, and fo generation of Poisson variates. 20. FORM 1473 EDITION OF INOV 66 IS OBSOLETE 			
 19. KEY WORDS (Continue on reverse elde II necessary and Identify by block number) Poisson processes, nonhomogeneous Poisson processes, one-dimensional non- homogeneous Poisson processes, two-dimensional nonhomogeneous Poisson pro- cesses, simulation of point processes, thinning, search and detection, acceptance-rejection, order statistics. 20. ABSTRACT (Continue on reverse elde II necessary and Identify by block number) A simple and relatively efficient method for simulating one-dimensional an two-dimensional nonhomogeneous Poisson processes is presented. The method is applicable for any rate function and is based on controlled deletion of points in a Poisson process whose rate function dominates the given rate function. In its simplest implementation, the method obviates the need fo numerical integration of the rate function, for ordering of points, and fo generation of Poisson variates. DD 1 JAN 73 1473 EDITION OF 1 NOV 65 IS OBSOLETE 			
 Poisson processes, nonhomogeneous Poisson processes, one-dimensional nonhomogeneous Poisson processes, two-dimensional nonhomogeneous Poisson processes, simulation of point processes, thinning, search and detection, acceptance-rejection, order statistics. Asstract (Continue on reverse elds if necessary and identify by block number) A simple and relatively efficient method for simulating one-dimensional an two-dimensional nonhomogeneous Poisson processes is presented. The method is applicable for any rate function and is based on controlled deletion of points in a Poisson process whose rate function dominates the given rate function. In its simplest implementation, the method obviates the need for numerical integration of the rate function, for ordering of points, and fo generation of Poisson variates. DD room 1473 EDITION OF INCOMENDATE 	19. KEY WORDS (Continue on reverse eide if necessary and identify by block numb		
 ADSTRACT (Continue on reverse side if necessary and identify by block number) A Simple and relatively efficient method for simulating one-dimensional an two-dimensional nonhomogeneous Poisson processes is presented. The method is applicable for any rate function and is based on controlled deletion of points in a Poisson process whose rate function dominates the given rate function. In its simplest implementation, the method obviates the need fo numerical integration of the rate function, for ordering of points, and fo generation of Poisson variates. 	Poisson processes, nonhomogeneous Poisson proce	esses, one-dimensional non-	
ABSTRACT (Continue on reverse elde 11 necessary and identify by block number) A simple and relatively efficient method for simulating one-dimensional an two-dimensional nonhomogeneous Poisson processes is presented. The method is applicable for any rate function and is based on controlled deletion of points in a Poisson process whose rate function dominates the given rate function. In its simplest implementation, the method obviates the need fo numerical integration of the rate function, for ordering of points, and fo generation of Poisson variates.	cesses, simulation of point processes, two-dimensional	nonnomogeneous roisson pro-	
ABSTRACT (Continue on reverse elde II necessary and identify by block number) A simple and relatively efficient method for simulating one-dimensional an two-dimensional nonhomogeneous Poisson processes is presented. The method is applicable for any rate function and is based on controlled deletion of points in a Poisson process whose rate function dominates the given rate function. In its simplest implementation, the method obviates the need fo numerical integration of the rate function, for ordering of points, and fo generation of Poisson variates.	acceptance-rejection, order statistics.	, coaren ana acceltion,	
A simple and relatively efficient method for simulating one-dimensional an two-dimensional nonhomogeneous Poisson processes is presented. The method is applicable for any rate function and is based on controlled deletion of points in a Poisson process whose rate function dominates the given rate function. In its simplest implementation, the method obviates the need fo numerical integration of the rate function, for ordering of points, and fo generation of Poisson variates.		ar)	
two-dimensional nonhomogeneous Poisson processes is presented. The method is applicable for any rate function and is based on controlled deletion of points in a Poisson process whose rate function dominates the given rate function. In its simplest implementation, the method obviates the need fo numerical integration of the rate function, for ordering of points, and fo generation of Poisson variates.	In the second se	wilating one-dimensional and	
is applicable for any rate function and is based on controlled deletion of points in a Poisson process whose rate function dominates the given rate function. In its simplest implementation, the method obviates the need fo numerical integration of the rate function, for ordering of points, and fo generation of Poisson variates.	ABSTRACT (Continue on reverse elde if necessary and identify by block numb A simple and relatively efficient method for st		
points in a Poisson process whose rate function dominates the given rate function. In its simplest implementation, the method obviates the need fo numerical integration of the rate function, for ordering of points, and fo generation of Poisson variates.	ABSTRACT (Continue on reverse eide it necessary and identify by block numb A simple and relatively efficient method for si two-dimensional nonhomogeneous Poisson processe	es is presented. The method	
DD FORM 1473 EDITION OF 1 NOV 65 IS OBSOLETE	ABSTRACT (Continue on reverse elde if necessary and identify by block numb A simple and relatively efficient method for si two-dimensional nonhomogeneous Poisson processe is applicable for any rate function and is base	es is presented. The method ed on controlled deletion of	
generation of Poisson variates.	ABSTRACT (Continue on reverse elde il necessory and identify by block numb A simple and relatively efficient method for si two-dimensional nonhomogeneous Poisson processe is applicable for any rate function and is base points in a Poisson process whose rate function	es is presented. The method ed on controlled deletion of a dominates the given rate	
DD 1 JAN 73 EDITION OF 1 NOV 65 IS OBSOLETE	ABSTRACT (Continue on reverse elde it necessary and identify by block numb A simple and relatively efficient method for si two-dimensional nonhomogeneous Poisson processe is applicable for any rate function and is base points in a Poisson process whose rate function function. In its simplest implementation, the	es is presented. The method ed on controlled deletion of a dominates the given rate method obviates the need fo	
DD 1 JAN 73 EDITION OF 1 NOV 65 IS OBSOLETE	A simple and relatively efficient method for sitt two-dimensional nonhomogeneous Poisson processes is applicable for any rate function and is base points in a Poisson process whose rate function function. In its simplest implementation, the numerical integration of the rate function, for generation of Poisson variates.	es is presented. The method ed on controlled deletion of a dominates the given rate method obviates the need fo c ordering of points, and fo	
	ABSTRACT (Continue on reverse elde it necessary and identify by block numb A simple and relatively efficient method for si two-dimensional nonhomogeneous Poisson processe is applicable for any rate function and is base points in a Poisson process whose rate function function. In its simplest implementation, the numerical integration of the rate function, for generation of Poisson variates.	es is presented. The method ed on controlled deletion of a dominates the given rate method obviates the need for c ordering of points, and for	

SIMULATION OF NONHOMOGENEOUS POISSON PROCESSES BY THINNING

by

P. A. W. Lewis^{*} Naval Postgraduate School Monterey, CA 93940

G. S. Shedler IBM Research Laboratory San Jose, CA 95193



JUNE 1978

^{*}Support under Office of Naval Research Grant NR-42-343 is gratefully acknowledged.

SIMULATION OF NONHOMOGENEOUS POISSON PROCESSES BY THINNING

by

P. A. W. Lewis Naval Postgraduate School Monterey, CA 93940

and

G. S. Shedler IBM Research Laboratory San Jose, CA 95193

ABSTRACT

A simple and relatively efficient method for simulating one-dimensional and two-dimensional nonhomogeneous Poisson processes is presented. The method is applicable for any rate function and is based on controlled deletion of points in a Poisson process whose rate function dominates the given rate function. In its simplest implementation, the method obviates the need for numerical integration of the rate function, for ordering of points, and for generation of Poisson variates.

KEYWORDS: Poisson processes, nonhomogeneous Poisson processes, one-dimensional nonhomogeneous Poisson processes, two-dimensional nonhomogeneous Poisson processes, simulation of point processes, thinning, search and detection, acceptance-rejection, order statistics.

1. INTRODUCTION

The one-dimensional nonhomogeneous Poisson process (see e.g., Cox and Lewis, 1966, pp. 28-29; Çinlar, 1975, pp. 94-101) has the characteristic properties that the numbers of points in any finite set of nonoverlapping intervals are mutually independent random variables, and that the number of points in any interval has a Poisson distribution. The most general nonhomogeneous Poisson process can be defined in terms of a monotone nondecreasing right-continuous function $\Lambda(\mathbf{x})$ which is bounded in any finite interval. Then the number of points in any finite interval, for example $(0, x_0]$, has a Poisson distribution with parameter $\mu_0 = \Lambda(x_0) - \Lambda(0)$. In this paper it is assumed that $\Lambda(x)$ is continuous, but not necessarily absolutely continuous. The right derivative $\lambda(\mathbf{x})$ of $\Lambda(\mathbf{x})$ is called the rate function of the process; $\Lambda(\mathbf{x})$ is called the integrated rate function and has the interpretation that for x > 0, $\Lambda(x) - \Lambda(0) = E[N(x)]$, where N(x)is the total number of points in (0,x]. Note that $\lambda(x)$ may jump at points at which $\Lambda(\mathbf{x})$ is not absolutely continuous. In contrast to the homogeneous Poisson process, i.e., $\lambda(x)$ a constant (usually denoted by λ), the intervals between the points in a one-dimensional nonhomogeneous Poisson process are neither independent nor identically distributed.

Applications of the one-dimensional nonhomogeneous Poisson process include modelling of the incidence of coal-mining disasters (Cox and Lewis, 1966), the arrivals at an intensive care unit (Lewis, 1972), transaction processing in a data base management system (Lewis and Shedler, 1976b), occurrences of major

freezes in Lake Constance (Steinijans, 1976), and geomagnetic reversal data (Reyment, 1976). The statistical analysis of trends in a one-dimensional nonhomogeneous Poisson process, based on the assumption of an exponential polynomial rate function, is discussed by Cox and Lewis (1966), Cox (1972), Lewis (1972), and Lewis and Shedler (1976b).

There are a number of methods for simulating nonhomogeneous Poisson process which we review briefly.

(i) Time-scale transformation of a homogeneous (rate one) Poisson process via the inverse of the (continuous) integrated rate function $\Lambda(\mathbf{x})$ constitutes a general method for generation of the nonhomogeneous Poisson process (cf., Çinlar, 1975 pp. 96-97). This method is based on the result that X_1, X_2, \ldots, T_n are the points in a nonhomogeneous Poisson process with continuous integrated rate function $\Lambda(x)$ if and only if $X_1^* = \Lambda(X_1)$, $X_2' = \Lambda(X_2)$, ..., are the points in a homogeneous Poisson process of rate one. The time-scale transformation method is a direct analogue of the inverse probability integral transformation method for generating (continuous) nonuniform random numbers. For many rate functions, inversion of $\Lambda(x)$ is not simple and must be done numerically; cf., Gilchrist (1977) and Patrow (1977). The resulting algorithm for generation of the nonhomogeneous Poisson process may be far less efficient than generation based on other methods; see e.g., Lewis and Shedler (1976a, 1977) and Patrow (1977) for discussion of special methods for efficiently generating the nonhomogeneous Poisson process with log-linear and log-quadratic rate functions.

- (ii) A second general method for generating a nonhomogeneous Poisson process with integrated rate function $\Lambda(\mathbf{x})$ is to generate the intervals between points individually, an approach which may seem more natural in the event-scheduling approach to simulation. Thus, given the points $X_1 = x_1, X_2 = x_2, \dots, X_i = x_i$, with $x_1 < x_2 < \cdots < x_i$, the interval to the next point, $x_{i+1} - x_i$, is independent of x_1, \ldots, x_{i-1} and has distribution function $F(x) = 1 - \exp[-\{\Lambda(x_i + x) - \Lambda(x_i)\}]$. It is possible to find the inverse distribution function $F^{-1}(\cdot)$, usually numerically, and generate $X_{i+1} - X_i$ according to $X_{i+1} - X_i = F^{-1}(U_i)$, where U_i is a uniform random number on the interval (0,1). Note, however, that this not only involves computing the inverse distribution function for each interval $X_{i+1} - X_i$, but that each distribution has different parameters and possibly a different form. An additional complication is that $X_{i+1} - X_i$ is not necessarily a proper random variable, i.e., there may be positive probability that $X_{i+1} - X_i$ is infinite. It is necessary to take this into account for each interval $X_{i+1} - X_i$ before the inverse probability integral transformation is applied. The method is therefore very inefficient with respect to speed, more so than the time-scale transformation method.
- (iii) In a third method, simulation of a nonhomogeneous Poisson process in a fixed interval $(0, x_0]$ can be reduced to the generation of a Poisson number of order statistics from a fixed density function by the following result (cf., Cox and Lewis, 1966, p. 45). If X_1, X_2, \ldots, X_n are the points of the

nonhomogeneous Poisson process in $(0, x_0]$, and if $N(x_0) = n$, then conditional on having observed n(> 0) points in $(0, x_0]$, the X_i are distributed as the order statistics from a sample of size n from the distribution function $\{\Lambda(x) - \Lambda(0)\}/\{\Lambda(x_0) - \Lambda(0)\}$, defined for $0 < x \leq x_0$. Generation of the nonhomogeneous Poisson process based on order statistics is in general more efficient (with respect to speed) than either of the previous two methods. Of course, a price is paid for this greater efficiency. First, it is necessary to be able to generate Poisson variates, and second, more memory is needed than in the interval-by-interval method in order to store the sequence of points. Enough memory must be provided so that with very high probability the random number of points generated in the interval can be stored. Recall that the number of points in the interval $(0, x_0]$ has a Poisson distribution with mean $\mu_0 = \Lambda(x_0) - \Lambda(0)$. Memory of size, e.g., $\mu_0 + 4\mu_0^{1/2}$ will ensure that overflow will occur on the average in only one out of approximately every 40,000 realizations. This probability is small enough so that in the case of overflow, the realization of the process can generally be discarded.

(iv) Again, there is a very particular and very efficient method for simulation of nonhomogeneous Poisson processes with log-linear rate function (Lewis and Shedler, 1976a) which, at the cost of programming complexity and memory, can be used to obtain an efficient simulation method for other rate functior ., as in Lewis and Shedler (1977).

In this paper a new method is given for simulating a nonhomogeneous Poisson process which is not only conceptually simple, but is also computationally simple and relatively efficient. In fact, at the cost of some efficiency, the method can be applied to simulate the given nonhomogeneous Poisson process <u>without the</u> <u>need for numerical integration or routines for generating Poisson</u> <u>variates</u>. Used in conjunction with the special methods given by Lewis and Shedler (1976a, 1977), the method can be used to generate quite efficiently nonhomogeneous Poisson processes with rather complex rate function, in particular combinations of long-term trends and fixed-cycle effects. The method is also easily extended to the problem of generating the two-dimensional nonhomogeneous Poisson process.

÷. ?'

2. SIMULATION OF ONE-DIMENSIONAL NONHOMOGENEOUS POISSON PROCESSES

Simulation of a nonhomogeneous Poisson process with general rate function $\lambda(\mathbf{x})$ in a fixed interval can be based on thinning of a nonhomogeneous Poisson process with rate function $\lambda^*(\mathbf{x}) \geq \lambda(\mathbf{x})$. The basic result is

THEOREM 1. Consider a one-dimensional nonhomogeneous Poisson process $\{N^{*}(x):x \ge 0\}$ with rate function $\lambda^{*}(x)$, so that the number of points, $N^{*}(x_{0})$, in a fixed interval $(0,x_{0}]$ has a Poisson distribution with parameter $\mu_{0}^{*} = \Lambda^{*}(x_{0}) - \Lambda^{*}(0)$. Let $x_{1}^{*}, x_{2}^{*}, \ldots, x_{N}^{*}_{*}(x_{0})$ be the points of the process in the interval

 $(0, \mathbf{x}_0]$. Suppose that for $0 \le \mathbf{x} \le \mathbf{x}_0$, $\lambda(\mathbf{x}) \le \lambda^*(\mathbf{x})$. For $\mathbf{i} = 1, 2, ..., n$, delete the point \mathbf{x}_1^* with probability $1 - \lambda(\mathbf{x}_1^*)/\lambda^*(\mathbf{x}_1^*)$; then the remaining points form a nonhomogeneous Poisson process $\{N(\mathbf{x}):\mathbf{x} \ge 0\}$ with rate function $\lambda(\mathbf{x})$ in the interval $(0, \mathbf{x}_0]$.

<u>Proof.</u> Since $\{N^{*}(x):x \ge 0\}$ is a nonhomogeneous Poisson process, and points are deleted independently, it is clear that the number of points in $\{N(x):x \ge 0\}$ in any set of nonoverlapping intervals are mutually independent random variables. Thus, it is sufficient to show that the number of points N(a,b) in $\{N(x);x \ge 0\}$ in an arbitrary interval (a,b] with $0 \le a < b \le x_0$ has a Poisson distribution with parameter $\Lambda(b) - \Lambda(a)$. Observe that with $p(a,b) = \{\Lambda(b) - \Lambda(a)\}/\{\Lambda^{*}(b) - \Lambda^{*}(a)\}$, we have the conditional probability

$$P\{N(a,b) = n | N^{*}(a,b)=k\} = \begin{cases} 1 & \text{if } n=k=0 \\ \{p(a,b)\}^{n}\{1-p(a,b)\}^{k-n} & \text{if } k \ge n \ge 0 \\ & \text{and } k \ge 1 \\ 0 & \text{if } n \ge 1 \\ & \text{and } k < n \\ \end{cases}$$
(1)

Equation (1) is a consequence of the well-known result that, conditional on n (>0) points in the interval (a,b], the joint density of the n points in the process $\{N^*(x):x \ge 0\}$ is

 $\lambda^*(\mathbf{x}_1) \cdots \lambda^*(\mathbf{x}_n) / \{\Lambda^*(\mathbf{b}) - \Lambda^*(\mathbf{a})\}^n$. The desired result is obtained in a straightforward manner from Equation (1) by removing the condition.

Theorem 1 is the basis for the method of simulating nonhomogeneous Poisson processes given in this paper.

ALGORITHM 1. One-dimensional nonhomogeneous Poisson process.

1. Generate points in the nonhomogeneous Poisson process $\{N^{*}(x)\}$ with rate function $\lambda^{*}(x)$ in the fixed interval $(0,x_{0}]$. If the number of points generated, n^{*} , is such that $n^{*} = 0$, exit; there are no points in the process $\{N(x)\}$.

2. Denote the (ordered) points by $x_1^*, x_2^*, \ldots, x_{n^*}^*$. Set i = 1 and k = 0.

- 3. Generate U_i , uniformly distributed between 0 and 1. If $U_i \leq \lambda(x_i^*) / \lambda^*(x_i^*)$, set k equal to k+1 and $X_k = x_i^*$.
- 4. Set i equal to i+1. If $i \leq n^*$, go to 3.
- 5. Return X_1, X_2, \ldots, X_n , where n = k, and also n.

In the case where

- (i) {N^{*}(x)} is a homogeneous Poisson process with $\lambda^*(x) = \lambda^*$;
- (ii) the minimum of $\lambda(\mathbf{x})$, say λ , is known, and
- (iii) generation of uniformly distributed variates is computationally costly,

considerable speedup can be obtained by noting that X_i^* is always accepted if $U_i \leq \lambda/\lambda^*$. This obviates, in some cases, computation

of $\lambda(\mathbf{x})$, which is the main source of inefficiency in the algorithm. Moreover, in this case $\lambda^* U_{\underline{i}} / \underline{\lambda}$ can be used as the next uniformly distributed variate.

The method of thinning in this simple form, i.e., $\lambda^*(\mathbf{x}) = \lambda^* \ge \max_{0 \le \mathbf{x} \le \mathbf{x}_0} \lambda(\mathbf{x})$, can also be used to provide an algorithm for generating a nonhomogeneous Poisson process on an interval-by-interval basis, as discussed in subsection (ii) of Section 1. The interval to the next point $\mathbf{x}_{i+1} - \mathbf{x}_i$ is obtained by generating and cumulating exponential (λ^*) random numbers $\mathbf{E}_1^*, \mathbf{E}_2^*, \ldots$, until for the first time $\mathbf{U}_j \le \lambda(\mathbf{x}_1 + \mathbf{E}_1^* + \cdots + \mathbf{E}_j^*)/\lambda^*$, where the \mathbf{U}_j are independent uniform random numbers between 0 and 1. This algorithm is considerably simpler than the interval-by-interval algorithm of Section 1 since it requires no numerical integration, only the availability of uniform random numbers.

3. DISCUSSION OF THE METHOD OF THINNING

(i) Relationship to acceptance-rejection method

The method of thinning of Algorithm 1 is essentially the obverse of the conditional method of Section 1, using conditioning and acceptance-rejection techniques to generate the random variables with density function $\lambda(\mathbf{x})/\{\Lambda(\mathbf{x}) - \Lambda(0)\}$ (Lewis and Shedler, 1977, Algorithm 3). The differences are subtle, but computationally important. In the acceptance-rejection method, it is first necessary to generate a Poisson variate with mean $\mu_0 = \Lambda(\mathbf{x}_0) - \Lambda(0)$, and this involves an integration of the rate function $\lambda(\mathbf{x})$. Then the Poisson (μ_0) number, n, of variates generated by acceptance-rejection must be ordered to give $\mathbf{x}_1, \mathbf{x}_2, \ldots, \mathbf{x}_n$.

(ii) Simplest form of the thinning algorithm

In the simplest form of the method of thinning, $\lambda^{*}(\mathbf{x})$ is taken to be a constant λ^{*} , so that, for instance, the points $\mathbf{x}_{1}^{*}, \mathbf{x}_{2}^{*}, \ldots, \mathbf{x}_{n^{*}}^{*}$ can be generated by cumulating exponential (λ^{*}) variates until the sum is greater than \mathbf{x}_{0} (cf., Lewis and , Shedler, 1976a, Algorithm 1). Thinning is then applied to the generated points. No ordering, no integration of $\lambda(\mathbf{x})$ and no generator of Poisson variates is required. Of course for both algorithms to be efficient, computation of $\lambda(\mathbf{x})$ and $\lambda^{*}(\mathbf{x})$ must be easy relative to computation of the inverse of $\Lambda(\mathbf{x})$.

(iii) Efficiency

For the thinning algorithm (as well as the algorithm based on conditioning and acceptance-rejection) efficiency, as measured by the number of points deleted, is proportional to $\mu_0/\mu_0^* =$ $\{\Lambda(\mathbf{x}_0) - \Lambda(0)\}/\{\Lambda^*(\mathbf{x}_0) - \Lambda^*(0)\}$; this is the ratio of the areas between 0 and \mathbf{x}_0 under $\lambda(\mathbf{x})$ and $\lambda^*(\mathbf{x})$. Thus, $\lambda^*(\mathbf{x})$ should be as close as possible to $\lambda(\mathbf{x})$ consistent with ease of generating the nonhomogeneous Poisson process $\{N^*(\mathbf{x}):\mathbf{x} \ge 0\}$.

(iv) An example: fixed cycle plus trend

To illustrate the applicability of the thinning algorithm, consider its use in conjunction with the algorithms given by Lewis and Shedler (1976a, 1977) for log-linear and log-quadratic rate functions. Assume that it is necessary to simulate a nonhomogeneous Poisson process whose rate function increases quadratically with time but also has a fixed-period cycle, e.g.,

 $\lambda(\mathbf{x}) = \exp\{\alpha_0 + \alpha_1 \mathbf{x} + \alpha_2 \mathbf{x}^2 + K \sin(\omega_0 \mathbf{x} + \theta)\},\$

 $0 \leq \mathbf{x} \leq \mathbf{x}_0; \quad \mathbf{K} \geq 0; \quad 0 < \theta \leq 2\pi; \quad \omega_0 > 0.$

This is the model found by Lewis (1972) for arrivals at an intensive care unit, where there is a strong time-of-day effect. Thus if $\omega_0 = 2\pi/T_0$, then the period $T_0 = 1$ day. Computation of $\Lambda^{-1}(\cdot)$ is difficult. To determine $\lambda^*(\mathbf{x})$, note that

$$\lambda(\mathbf{x}) \leq \lambda^{*}(\mathbf{x}) = \exp\{\alpha_{0} + \mathbf{K} + \alpha_{1}\mathbf{x} + \alpha_{2}\mathbf{x}^{2}\},$$
10

and therefore

$$\lambda(\mathbf{x})/\lambda(\mathbf{x}) = \exp[K\{1 - \sin(\omega_0 \mathbf{x} + \theta)\}]$$
.

Thus in Step 3 of Algorithm 1, U_i is compared to $\exp[K\{1 - \sin(\omega_0 X_i^{*} + \theta)\}]$. Equivalently, if unit exponential variates E_i are available, it is faster to compare E_i to $K\{1 - \sin(\omega_0 X_i^{*} + \theta)\}$, accepting X_i^{*} if $E_i > K\{1 - \sin(\omega_0 X_i^{*} + \theta)\}$.

The main computational expense here is generation of the E_i and computation of the sine function, both n^* times. The expense involved in computation of the sine function can be reduced by noting that the point X_i^* is always accepted if E_i is greater than 2K. This will be a great saving if the cyclic effect is minor (K small). The number of E_i generated can be reduced by noting that if, in one step of the algorithm, E_i is observed to be greater than δ , then $E_i^* = E_i - \delta$ can be used as an (independent) unit exponential variate in the next step. The above procedure can be extended to the case of a trend with two fixed-period cycles, e.g., a time-of-day and a time-of-week effect.

4. SIMULATION OF TWO-DIMENSIONAL HOMOGENEOUS POISSON PROCESSES

The two-dimensional homogeneous Poisson process (of rate $\lambda > 0$) is defined by the properties that the numbers of points in any finite set of nonoverlapping regions having areas in the usual geometric sense are mutually independent, and that the number of points in any region of area A has a Poisson distribution with mean λA ;

see, e.g., Karlin and Taylor (1975), pp. 31-32. Note that the number of points in a region R depends on its area, but not on its shape or location. The homogeneous Poisson process arises as a limiting two-dimensional point process with respect to a number of limiting operations; cf., Goldman (1967a,b). Properties of the process are given by Miles (1970). Applications of the twodimensional homogeneous Poisson process to problems in ecology and forestry have been discussed by Thompson (1955) and Holgate (1972). The model also arises in connection with naval search and detection problems.

In considering the two-dimensional homogeneous Poisson process, projection properties of the process depend quite critically on the geometry of the regions considered. These projection properties are simple for rectangular and circular regions, and make simulation of the homogeneous process quite easy. We consider these two cases separately.

(i) Homogeneous Poisson Processes in a Rectangle

The following two theorems form the basis for simulation of the two-dimensional homogeneous Poisson process in a rectangle.

<u>THEOREM 2</u>. Consider a two-dimensional homogeneous Poisson process of rate λ , so that the number of points in a fixed rectangle $R = \{(x,y): 0 < x \leq x_0, 0 < y \leq y_0\}$ has a Poisson distribution with parameter $\lambda x_0 y_0$. If $(X_1, Y_1), (X_2, Y_2), \dots, (X_N, Y_N)$ denote the position of the points of the process in R, labelled so that

 $x_1 < x_2 < \cdots < x_N$, then x_1, x_2, \ldots, x_N form a one-dimensional homogeneous Poisson process on $0 < x \le x_0$ of rate λy_0 . If the points are relabelled $(x_1', y_1'), (x_2', y_2'), \ldots, (x_N', y_N')$ so that $y_1' < y_2' < \cdots < y_N'$, then y_1', y_2', \ldots, y_N' form a one-dimensional homogeneous Poisson process on $0 < y \le y_0$ of rate λx_0 .

<u>Proof</u>. The number of points in an interval on the x-axis, say, (a,b] is the number of points in the rectangle bounded by the lines x = a, x = b, y = 0, and $y = y_0$. This number is therefore independent of the number of points in any similar nonoverlapping rectangle bounded on the x-axis by x = a', x = b', i.e., the number of points in the interval (a',b']. This establishes the independent increment property for a one-dimensional Poisson process. The Poisson distribution of the number of points in (a,b) follows from the fact that it is equal to the number of points in the rectangle bounded by x = a, x = b, y = 0, and $y = y_0$, and the latter has a Poisson distribution with parameter λy_0 (b-a). An analogous argument shows that the process formed on the y-axis by Y'_1 , Y'_2 , ..., Y'_N is Poisson.

Conditional properties of the Poisson process in a rectangle are established next. The important thing to note is that while the processes obtained by projection of the points onto the x and y axes are not independent, there is a type of conditional independence which makes it easy to simulate the two-dimensional process.

<u>THEOREM 3</u>. Assume that a two-dimensional homogeneous Poisson process of rate λ is observed in a fixed rectangle $R = \{(x,y): 0 \le x \le x_0, 0 \le y \le y_0\}$, so that the number of points in R, N(R), has a Poisson distribution with parameter $\lambda x_0 y_0$. If N(R) = n > 0 and if $(X_1, Y_1), (X_2, Y_2), \dots, (X_n, Y_n)$ denote the points, labelled so that $X_1 \le X_2 \le \dots \le X_n$, then conditional on having observed n points in R, the X_1, X_2, \dots, X_n are uniform order statistics on $0 \le x \le x_0$, and Y_1, Y_2, \dots, Y_n are independent and uniformly distributed on $0 \le y \le y_0$, independent of the X_i .

Proof. If there are N points in the rectangle, form N vertical strips, from 0 to y_0 and from X_i to $X_i + dx_i$, such that each strip contains only one of the N points. The position of Y_i on the vertical line through X_i is that of an event in a Poisson process of rate λdx_i , given that only one event occurs. But this means that Y, is uniformly distributed between 0 and y_0 . Moreover, this is true irrespective of where X_i occurs; therefore Y_i is independent of X_i . Also, occurrences in all N strips are independent, and therefore Y_{i} is independent of the other Y_{i} and X_{i} positions, $j \neq i$. Thus the Y_i are a random sample of size N from a uniform $(0, y_0)$ distribution, independent of the X_i . Now condition on N = n (> 0); since by Theorem 2 the X_i form a Poisson process they are, by well-known results, order statistics from a uniform $(0, x_0)$ sample and are independent of the fixed size Y_i population; thus the pairs (X_i, Y_i) are mutually independent.

<u>COROLLARY</u>. Denote the Poisson points by (X_1, Y_1) , (X_2, Y_2) , ..., where the index does not necessarily denote an ordering on either axis. Conditionally, the pairs (X_1, Y_1) , ..., (X_N, Y_N) are independent random variables. Furthermore, for each pair, X_i is distributed uniformly between 0 and x_0 , independently of Y_i , which is uniformly distributed between 0 and y_0 .

From the two theorems, the following simulation procedure is obtained.

ALGORITHM 2. Two-dimensional homogeneous Poisson process in a rectangle.

- 1. Generate points in the one-dimensional homogeneous Poisson process of rate λy_0 on $(0, x_0]$. If the number of points generated, n, is such that n = 0, exit; there are no points in the rectangle.
- 2. Denote the points generated by $\lambda_1 < X_2 < \cdots < X_n$.
- 3. Generate Y_1, Y_2, \ldots, Y_n as independent, uniformly distributed random numbers on $(0, y_0]$.
- 4. Return (X_1, Y_1) , (X_2, Y_2) , ..., (X_n, Y_n) as the coordinates of the two-dimensional homogeneous Poisson process in the rectangle, and n.

Note that generation of the points X_1, X_2, \ldots, X_N in Steps 1 and 2 can be accomplished by cumulating exponential (λy_0) random numbers. Alternatively, after generating a Poisson random number N = n (with parameter $\lambda x_0 y_0$), n independent, uniformly distributed random numbers on $(0, x_0]$ can be ordered; see Lewis and Shedler (1976a), p. 502.

Another algorithm for generation of the two-dimensional Poisson process in a rectangle can be based on the Corollary to Theorem 3.

(ii) Homogeneous Poisson Processes in a Circle

The following two theorems form the basis for simulation of the two-dimensional homogeneous Poisson process in a fixed circle of radius r_0 .

Fix the origin and initial line of polar coordinates r and θ so that the origin is the center of the circle and the initial line is horizontal. We consider the projection of the points (R_i, θ_i) , of the Poisson process circularly onto the r-axis (R_i) and radially onto the circumferential θ -axis (θ_i) . The number of points projected onto the r-axis in the interval (0,r], where $r \leq r_0$, is the number of points in the circle of radius r and area πr^2 ; thus the number of points in (0,r]has a Poisson distribution with parameter $\lambda \pi r^2$. Consequently, if the projection process on the r-axis is a Poisson process, it must have integrated rate function $\Lambda(r) = \lambda \pi r^2$, with $\Lambda(0) = 0$.

Similarly, the number of points on the circumferential arc of the fixed circle (radius r_0) from 0 to θ is the number of points in the sector of the circle defined by radial lines at angles 0 and θ ; thus, the number of points on the arc from 0 to θ has a Poisson distribution with parameter $\lambda \pi r_0^2 \times \frac{\theta}{2\pi}$ = $\theta \lambda r_0^2/2$. Accordingly, if the projection process on the θ -axis is a Poisson process, it must have integrated rate function $\Lambda(\theta) = \theta \lambda r_0^2/2$, with $\Lambda(0) = 0$.

We now assert that the projection processes are in fact Poisson processes. Since proofs of these theorems are directly analogous to the proofs of Theorems 2 and 3, they are omitted.

<u>THEOREM 4</u>. Consider a two-dimensional homogeneous Poisson process of rate λ so that the number N of points in a fixed circular area C of radius r_0 and area πr_0^2 has a Poisson distribution with parameter $\lambda \pi r_0^2$. If (R_1, θ_1) , (R_2, θ_2) , ..., (R_N, θ_N) denote the points of the process in C, labelled so that $R_1 \leq R_2 \leq \cdots \leq R_N$, then R_1, R_2, \ldots, R_N form a one-dimensional nonhomogeneous Poisson process on $0 \leq r \leq r_0$ with rate function $\lambda(r) = 2\pi\lambda r$. If the points are relabelled (R_1^i, θ_1^i) , (R_2^i, θ_2^i) , ..., (R_N^i, θ_N^i) so that $\theta_1^i \leq \theta_2^i \leq \cdots \leq \theta_N^i$, then $\theta_1^i, \theta_2^i, \ldots, \theta_N^i$ form a one-dimensional homogeneous Poisson process on $0 \leq \theta \leq 2\pi$ of rate $\lambda r_0^2/2$.

<u>THEOREM 5.</u> Assume that a two-dimensional Poisson process of rate λ is observed in a fixed circular area C of radius r_0 so that the number of points in C, N(C), has a Poisson distribution with parameter $\lambda \pi r_0^2$. If N(C) = n > 0 and if $(R_1, \theta_1), (R_2, \theta_2), \ldots, (R_n, \theta_n)$ with $R_1 \leq R_2 \leq \cdots \leq R_n$ denote the points, then conditional on having observed n points in C, the R_1, R_2, \ldots, R_n are order statistics from the density $f(r) = 2r/r_0^2$ concentrated on $0 \leq r \leq r_0$, and $\theta_1, \theta_2, \ldots, \theta_n$ are independent and uniformly distributed on $0 < \theta \leq 2\pi$, independent of the R_i .

These theorems lead to the following simulation procedure.

ALGORITHM 3. Two-dimensional homogeneous Poisson process in a circular area.

- 1. Generate n as a Poisson random number with parameter $\lambda \pi r_0^2$. If n = 0, exit; there are no points in C.
- 2. Generate n independent random numbers having density function $f(r) = 2r/r_0^2$ and order to obtain $R_1 < R_2 < \cdots < R_n$.
- 3. Generate $\theta_1, \theta_2, \dots, \theta_n$ independent, uniformly distributed random numbers on $(0, 2\pi]$.

4. Return
$$(R_1, \theta_1)$$
, (R_2, θ_2) , ..., (R_n, θ_n) , and n.

Note that the wedge-shaped density $2r/r_0^2$ can be generated by scaling the maximum of two independent uniform (0,1) random numbers.

Direct generation of homogeneous Poisson points in non-circular or non-rectangular regions is difficult. The processes obtained by projection of the points on the two axes are nonhomogeneous Poisson processes with complex rate functions determined by the geometry of the region. However, the conditional independence which is found in circular and rectangular regions (Theorems 3 and 5) for the processes on the two axes is not present. In particular, given that there are n points $(x_1, y_1), \ldots, (x_n, y_n)$ in a non-rectangular region, the pairs (x_i, y_i) are mutually independent, but x_i is in general not independent of y_i , $i = 1, \ldots, n$. Therefore, it is simpler to enclose the

region in either a circle or a rectangle, generate a homogeneous Poisson process in the enlarged area, and subsequently exclude points outside of the given region.

5. SIMULATION OF TWO-DIMENSIONAL NONHOMOGENEOUS POISSON PROCESSES

The two-dimensional nonhomogeneous Poisson process $\{N(x,y):x \ge 0, y \ge 0\}$ is specified by a positive rate function $\lambda(x,y)$ which for simplicity is assumed here to be continuous. Then the process has the characteristic properties that the numbers of points in any finite set of nonoverlapping regions having areas in the usual geometric sense are mutually independent, and that the number of points in any such region R has a Poisson distribution with mean $\Lambda(R)$; here $\Lambda(R)$ denotes the integral of $\lambda(x,y)$ over R, i.e., over the entire area of R.

Applications of the two-dimensional nonhomogeneous Poisson process include problems in forestry and naval search and detection. The use of the process as a model for the pattern of access to the storage subsystem of a computer system will be reported elsewhere. Detection and statistical analysis of trends in the two-dimensional nonhomogeneous Poisson process is discussed by Rantschler (1973).

Theorem 1 dealing with thinning of one-dimensional nonhomogeneous Poisson processes generalizes to two-dimensional nonhomogeneous Poisson processes. Thus, suppose that $\lambda(\mathbf{x},\mathbf{y}) \leq \lambda^{*}(\mathbf{x},\mathbf{y})$ in a fixed rectangular region of the plane. If a nonhomogeneous Poisson process with rate function $\lambda^{*}(\mathbf{x},\mathbf{y})$ is thinned according to $\lambda(\mathbf{x},\mathbf{y})/\lambda^{*}(\mathbf{x},\mathbf{y})$ (i.e., each point

 (X_{i}, Y_{i}) is deleted independently if a uniform (0,1) random number U_{i} is greater than $\lambda(X_{i}, Y_{i})/\lambda^{*}(X_{i}, Y_{i}))$, the result is a nonhomogeneous Poisson process with rate function $\lambda(x, y)$. The proof is a direct analogue of the proof for the one-dimensional case.

The nonhomogeneous Poisson process with rate function $\lambda(x,y)$ in an arbitrary but fixed region R can be generated by enclosing the region R either in a rectangle or a circle, and applying Algorithm 2 or Algorithm 3. The following procedure assumes that the region R has been enclosed in a rectangle R^{*}, and that $\lambda^* = \max\{\lambda(x,y): x, y \in R\}$ has been determined; here the bounding process is homogeneous with rate λ^* in the rectangle R^{*}.

ALGORITHM 4. Two-dimensional nonhomogenous Poisson process.

- 1. Using Algorithm 2, generate points in the homogeneous Poisson process of rate λ^* in the rectangle R^{*}. If the number of points, n^{*}, is such that n^{*} = 0, exit; there are no points in the nonhomogeneous Poisson process.
- 2. From the n^* points generated in 1, delete the points that are not in R, and denote the remaining points by $(x_1^*, y_1^*), (x_2^*, y_2^*), \dots, (x_m^*, y_m^*)$ with $x_1^* < x_2^* < \dots < x_m^*$. Set i = 1 and k = 0.
- 3. Generate U_i uniformly distributed between 0 and 1. If $U_i \leq \lambda (X_i^*, Y_i) / \lambda^*$, set k = k+1, $X_k = X_i^*$ and $Y_k = Y_i^*$.
- 4. Set i equal to i+1. If $i \leq m^*$, go to 3.
- 5. Return (X_1, Y_1) , (X_2, Y_2) , ..., (X_n, Y_n) , where n = k, and n.

It is not necessary that the bounding process have a constant rate λ^* . Theroems 2 and 4 can be generalized to certain cases where the process is nonhomogeneous (cf., Bartlett, 1974), for instance $\lambda(\mathbf{x}, \mathbf{y}) = \rho(\mathbf{x}) \psi(\mathbf{y})$. Thus, a tighter bounding process which is nonhomogeneous may possibly be obtained. It is not simple to see how much efficiency could be gained by doing this, as opposed to using a two-dimensional homogeneous Poisson process for the bounding process. Again, as in the one-dimensional case, savings in computing $\lambda(\mathbf{x}, \mathbf{y})$ can be obtained by computing its minimum beforehand, and the U_i 's can be reused by scaling.

6. COMPARISONS AND CONCLUDING REMARKS

The method of thinning presented in this paper for simulating one-dimensional and two-dimensional nonhomogeneous Poisson processes with given rate function can be carried out in a computationally simple way by using a bounding process which is homogeneous with a rate function equal to the maximum value of the given rate function. No numerical integration, ordering or generation of Poisson variates is required, only the ability to evaluate the given rate function. The thinning algorithm appears to be particularly attractive in the two-dimensional case where there seem to be no competing algorithms.

The thinning algorithm can also be implemented more efficiently at the cost of programming complexity and by using a nonhomogeneous bounding process. In particular the method can be used in conjunction with the special algorithms given by Lewis and Shedler (1976a, 1977).

It is also possible to extend the method of thinning to simulation of doubly stochastic or conditioned Poisson processes. This will be discussed elsewhere.

REFERENCES

- Ahrens, J. and Dieter, U. (1974a). Computer methods for sampling from gamma, beta, Poisson and binomial distributions. Computing 12, 223-46.
- Ahrens, J. and Dieter, J. (1974b). Non Uniform Random Numbers. Graz, Austria: Technische Hochschule in Graz.
- Bartlett, M.S. (1964). The spectral analysis of two-dimensional point processes. Biometrika 51, 299-311.
- Cinlar, E. (1975). Introduction to Stochastic Processes. Englewood Cliffs, New Jersey: Prentice-Hall.
- Cox, D.R. The statistical analysis of dependencies in point processes. In <u>Stochastic Point Processes</u>, Ed. P.A.W. Lewis, pp. 55-56. New York: Wiley.
- Cox, D.R. and Lewis, P.A.W. (1966). The Statistical Analysis of Series of Events. London: Methuen.
- Gilchrist, R. (1977). Some problems encountered in generating realisations from a useful non-homogeneous Poisson process. In Proceedings of the Eruopean Congress of Statisticians 1976, Ed. J. R. Barra et al. Amsterdam: North Holland.
- Goldman, J.R. (1967a). Stochastic point processes: limit theorems. Ann. Math. Stat. 38, 771-79.
- Goldman, J.R. (1967b). Infinitely divisible point processes in Rⁿ. J. Math. Anal. Appl. 17, 133-46.
- Holgate, P. (1972). The use of distance methods for the analysis of spatial distributions of points. In <u>Stochastic Point</u> Processes, Ed. P.A.W. Lewis, pp. 122-35. New York: Wiley.
- Karlin, S. and Taylor, H.M. (1975). A First Course in Stochastic Processes. New York: Academic Press.
- Lewis, P.A.W. (1972). Recent advances in the statistical analysis of univariate point processes. In Stochastic Point Processes, Ed. P.A.W. Lewis, pp. 1-54. New York: Wiley.
- Lewis, P.A.W. and Shedler, G.S. (1976a). Simulation of nonhomogeneous processes with log-linear rate function. Biometrika 63, 501-05.

- Lewis, P.A.W. and Shedler, G.S. (1976b). Statistical analysis of a non-stationary series of events in a data base system. IBM J. Res. Devel. 20, 465-82.
- Lewis, P.A.W. and Shedler, G.S. (1977). Simulation of nonhomogeneous Poisson processes with degree-two exponential polynomial rate function. IBM Research Report RJ 1953. San Jose, California.
- Miles, R.E. (1970). On the homogeneous planar Poisson point process. Mathematical Biosciences 6, 85-127.
- Patrow, M.L. (1977). A comparison of two algorithms for the simulation of non-homogeneous Poisson processes with degree-two exponential polynomial intensity function. M.S. Thesis., Dept. of Operations Research, Naval Postgraduate School. Monterey, California (National Technical Information Service, Springfield, Va.; AD-A047164).
- Rantschler, R.D. (1973). The detection and estimation of trends in two-dimensional Poisson processes. M.S. Thesis, Dept. of Operations Research, Naval Postgraduate School, Monterey, California (National Technical Information Service, Springfield, Va.; AD-A76136).
- Reyment, R.A. (1976). Geomagnetic reversal data statistically appraised. Tectonophysics 31, 73-91.
- Steinijans, V. (1976). A stochastic point process model of the occurrence of major freezes in Lake Constance. <u>Appl.</u> <u>Statistics 25, 58-61.</u>
- Thompson, H.R. (1955). Spatial point processes with applications to ecology. Biometrika 42, 102-15.
- Vere-Jones, D. (1970). Stochastic models for earthquake occurrence. J. Roy. Statist. Soc. B 32, 1-62.

23

ŧ

,

.

	_	NO.	OF	COPIES
STATISTICS AND FREBABILITY CFFICE OF NAVAL RESEARCH CODE 436 ARLINGION	FROGRAM		,	
CEEICE OF NAVAL RESEARCH	· · ·		-	
715 BROACHAY - 5TH FLOOR ATTN: DR. RJEER GRAFTON NEW YORK, NY	10003		1	
CIRECTCR OFFICE OF NAVAL RESEARCH BE 536 SCUTH CLARK STREET ATTN: DEPUTY AND CHIEF SCI	RANCH OFF			
LIBRARY			1	
SAN DIEGC CA	92152		1	
NAVY LIBPARY NATIONAL SPACE TECHNOLCGY L ATTN: NAVY LIBRARIAN	. A B			
MS	39522		1	
NAVAL ELECTRONIC SYSTEMS CO	SMMA ND			
ARLINGTON VA	20360	p.	1	
CIPECTOR NAVAL REAFARCH LAG	ORATORY			
COCE 2C29 WASHINGTON, C.C.	20375 .		1	
DEFENSE COCUMENTATION CENTS CAPEFON STATION	R		1	
ALEXANDRIA VIFGINIA 22314				
TECHNICAL INFORMATION DIVIS	SION .		1	
WASHINGTON, C. C.				

20375

۰,

CFFICE OF NAVAL RESEARCH SAN FRANCISCC AREA OFFICE 760 Market Street SAN FRANCISCL CALIFORNIA 94102	NO. OF	COPIES
TECHNICAL LIBRARY NAVAL DRENANCE STATICN	_	
INDIAN HEAE MARYLAND 20640	1	
NAVAL SHIP ENGINEERING CENTER PHILADELFHIA DIVISION TECHNICAL LIBRARY PHILADELPHIA PENNSYLVANIA 19112	1	
BUREAU CF NAVAL PRESONNEL DEPARTMENT OF THE NAVY TECHNICAL LIERARY WASHINGTON C. C. 20370	1	
LIERARY CODE 2212 NAVAL POSTGRACLATE SCHOOL MENTEREY CALIFORNIA 53940	1	
PRCF. M. AEDEL-HAMEED DEFARTMENT OF MATHEMATICS UNIVERSITY CF NORTH CARCLINA CHARLOTTE NC 28223	1	
PROF. T. W. ANCERSON DEFARTMENT OF STATISTICS STANFORC UNIVERSITY	.	
FROF. F. J. ANSCOMBE	1	
TALE LNIVERSITY NEW FAVEN CONNECTIOUT 06520	1	
PROF. L. A. AFCIAN INSITILTE OF INCUSTRIAL AD MINISTRATION UNION COLLEGE SCHENECTADY . NEW YORK 12308	1	
PROF. C. R. BAKER DEFARTMENT OF STATISTICS UNIVERSITY OF NOTRH CARGLINA CHAPEL FILL, NORTH CARCLINA		
27514	1	

PROF R. E. BECHOFER DEFARTMENT OF OPERATIONS RESEARCH CCRNELL UNIVERSITY ITHACA NEW YORK 1485C	1
FRCF. N. J. BERSHAD SCHOOL CF ENGINEERING UNIVERSITY OF CALIFORNIA IRVINE . CALIFORNIA S2664	1
P. J. BICKEL CEPARTMENT OF STATISTICS UNIVERSITY OF CALIFORNIA REFERENCE. STATISTICS	1
PROF L W RICCK	· •
DE FARTMENT OF WATHENATICS UNIVERSITY OF FITTSBURGH PITTSBURGH	T
15260	_
FREF. JOSEPH BLUM DEFT. OF MATHEMATICS, STATISTICS AND COMPUTER SCIENCE THE AMERICAN UNIVERSITY	1
ASHINGTON 20016	
PFOF. R. A. BRADLEY DEPARTMENT OF STATISTICS FLORIDA STATE UNIVERSITY	1
TALLAHASSEE , FLORIDA 32306	
PROF. R. E. BARLOW CFERATIONS RESEARCH CENTER COLLEGE OF ENGINEERING UNIVERSITY OF CALIFORNIA BEFKLEY CALIFORNIA 54720	1
MR. C. N BENNETT NAVAL COASTAL SYSTEMS LABCFATORY CODE P761 PANAMA CITY, FLCRIDA 32401	1
PROF. U. N. BHAT CCMPUTER SCIENCE / OPERATIONS RESEARCH CENTER SOUTHERN METHODIST UNIVERSITY CALLAS TEXAS 75275	1
PRCF. W. R. BLISCHKE DEPT. OF QUANTITATIVE BUSINESS ANALYSIS UNIVERSITY OF SOUTHERN CALIFOFNIA LOS ANGELES, CALIFORNIA	1

<u> 50007</u>

	DR. DERRILL J. BORDELGN NAVAL UNDERWATER SYSTEMS CENTER COCE 21 Newport	NO. OF COPIES
	RI C2 84 0	
	J. E. BOYER JR DEPT. OF STATISTICS SCUTHERN METHOLIST UNIVERSITY DALLAS TX 75275	1
	DR. J. CHANCRA U. S. ARMY RESEARCH P. G. BCX 12211 RESEARCH TRIANGLE PARK, NCFTH CAROLINA 27706	I
¥	PROF. F. CHERNOFF DEPT. OF MATHEMATICS MASS INSTITUTE OF TECHNOLOGY CAMBRINGE MASSACHUSETTS 02139	. 1
	PROF. C. DERMAN DEFARTMENT OF CIVIL ENGINEERING AND ENGINEERING MECHANICS COLUMBIA UNIVERSITY NEW YORK NEW YORK 10027	1
	FRCF. R. L. CISNEY VIRGINIA PCLYTECHNIC INSTITUTE AND STATE UNIVERSITY DEPT. CF INDLSTPIAL ENGINEERING AND CPERATIONS RESEARCH BLACKSBUFG, VA 24061	1
	FR. J. DOWLING DEFENSE LOGISTICS STUDIES INFORMATION EXCHANGE ARMY LOGISTICS MANAGEMENT CENTER FORT LEE VIRGINIA 20390	1
	PRCF J. D. ESARY CEPT. CF OFERATIONS RESEARCH AND ADMINISTRATIVE SCIENCES NA VAL POSTGRADUATE SCHOOL MONTEREY CALIFORNIA 53540	1
	DR. M. J. FISCHER DEFENSE COMMUNICATIONS AGENCY 1860 WIEHLE AVENUE RSTON VIRGINIA 22070	1

·

.

.

•

.

the training

PROF. C. P. GAVER DEFT. OF OPERATIONS RESEARCH NAVAL FESTERADLATE SCHEEL MONTEREY CA 53940	1
MR. GENE F. GLEISSNER APPLIED MATHEMATICS LAEORATORY DAVID TAYLCE NAVAL SHIP RESEARCH AND DEVELOPMENT CENTER BETHESCA MD 20084	1
PROF. S. S. GUPTA DEPARTMENT OF STATISTICS FURDUE UNIVERSITY LAFAYETTE INDIANA 47907	1
PROF. D. L. HANSON DEFT OF MATH. SCIENCES STATE UNIVERSITY OF NEW YORK, EINGHAMTON BINGHAMTON NY 13901	ŗ
PROF. F. J. HARRIS DEFT. OF ELECTRICAL ENGINEERING SAN CIEGC STATE UNIVERSITY SAN DIEGC CA 52182	1
FRCF. L. H. HERBACH DEPT. CF CFERATICNS RESEARCH AND SYSTEMS ANALYSIS POLYTECHNIC INSTITUTE CF NEW YORK BRCOKLYN NY 11201	1
PRCF. V. J. HINICH DEPARTMENT OF ECONOMICS VIRGINIA POLYTECHNIC INSTITUTE ANC STATE UNIVERSITY BLACKSEURG VIRGINIA 24C61	1
PROF. W. M. HIRSCH INSTITUTE OF MATHEMATICAL SCIENCES New York UNIVERSITY New York New York 10453	1
PROF. C. L. IGLEHART DEFARTMENT OF OPERATIONS RESEARCH STANFORD UNIVERSITY STANFORD. CALIFORNIA 94350	1

y

FRCF. J. E. KACANE DEPARTMENT CF STATISTICS CARNEGIE-MELLCN FITTSBURGH . FENNSYLVANIA 15213		ľ
DR. RICHARE LAU DIRECTOR CFFICE OF NAVAL RESEARCH BRANCH OFF 1030 EAST GREEN STREET FASADENA CA 91101		1
DR . A . R . LAUFER DIRECTOR OFFICE OF NAVAL RESEARCH ERANCH OFF 1030 EAST GREEN STREET PASADENA CA 91101		1
FROF. M. LEACEETTER DEPARTMENT OF STATISTICS UNIVERSITY OF NORTH CAROLINA CHAPEL HILL NORTH CAROLINA 27514		1
CR. J. S. LEE J. S. LEE ASSCCIATES, INC. 2001 JEFFERSON DAVIS HIGHWAY SUITE EC2 ARLINGTON VA 22202	•	1
PROF. L. C. LEE DE PARTMENT OF STATISTICS VIRGINIA POLYTECHNIC INSTITUTE AND STATE UNIVERSITY BLACK SEURG VA 24061		1
PROF. R. S. LEVENWORTH DEPT. OF INCUSTRIAL AND SYSTEMS ENGINEERING UNIVERSITY OF FLORIDA GAINSVILLE, FLORIDA 32611	•	1
PRCF G. LIEPERMAN Stanford Iniversity Department of cperations research Stanford California 94305	×	1

¥

ì

DR. JAMES F. MAAR National Security Agency Fort Meade , Maryland 20755
PROF. R. W. WACSEN DEFARTMENT OF STATISTICS UNIVERSITY OF MISSOURI COLUMEIA MC 65201
DR. N. R. MANN SCIENCE CENTER RECKWELL INTERNATIONAL CORPORATION F.C. BEX ICE5 THEUSAND CAKS CALIFORNIA 91360
DR . W. H. MARLOW PREGRAM IN LOGISTIES THE GEERGE WASHINGTEN UNIVERSITY 707 22ND STREET , N. W. WASHINGTON , C. C. 20037
PROF. E. MASRY DEFT. APPLIEC PHYSICS AND INFERMATION SERVICE UNIVERSITY OF CALIFORNIA LA JOLLA CALIFORNIA 52093
DF. BRUCE J. MCCONALD SCIENTIFIC DIRECTOR SCIENTIFIC LIAISON GROUF CFFICE OF NAVAL RESEARCH AMERICAN EMBASSY - TOKYO AFC SAN FRANCISCO 96503
FRCF. J. A. MUCKSTADT CEPT. CF CFERATIONS RESEARCH CCRNELL UNIVERSITY ITHACA, NEW YORK 19850
DF. JANET M. MYHRE THE INSTITUTE OF DECISION SCIENCE FOR BUSINESS AND PUBLIC FOLICY CLAREMONT MEN'S COLLEGE CLAREMONT CA 91711
MR. H. NISSELSCN BLREAU OF THE CENSUS PCCM 2C25 FRCERAL EVILCING 3 WASHINGTON , C. C. 2C33

ŧ

VISS B. S. ORLEANS NAVAL SEA SYSTEMS COMMAND (SEA 03F) RM 10508 ARLINGTON VIRGINIA 20360 1 FRGF. C. B OWEN CEPARTMENT OF STATISTICS SCUTHERN METHOCIST UNIVERSITY DALLAS TEXAS 75222 1 DR. A. PETRASOVITS ROOM 2078, FOCD AND DRUG ELDG. TUNNEY'S PASTURE OTTOWA, CNTARIC K1A-OL2, 1 CANADA 14 PROF. S. L. PHCENIX SIELEY SCHOOL OF MECHANICAL AND AEROSPACE ENGINEERING CCFNELL UNIVERSITY ITHACA NY 14850 1 14850 DR. A. L. POWELL CIRECTOR CFFICE OF NAVAL RESEARCH BRANCH OFF 495 SUMMER STREET BOSTON MA . C2210 1 MR. F. R. PRIORI CODE 224 CFERATIONSL TEST AND ONRS EVALLATION FORCE (OPTEVFOR) NORFOLK , VIFGINIA 20360 1 PROF. F. L. PUFI DEFT. CF MATHEMATICS F.C. BOX F INCIANA UNIVERSITY FOUNCATION BLOOMING TON IN 474 1 47401 PROF. H RCBBINS DEFARTMENT OF MATHEMATICS COLUMBIA UNIVERSITY NEW YORK, NEW YORK, ICO27 1

•

t

	FROF. M ROSENBLATT DEPARTMENT OF MATHEMATICS UNIVERSITY OF CALIFORNIA SAN DIEGO LA JOLLA CALIFORNIA \$2093	1
	FROF. S. M. ROSS College of Engineering University of California Befkeley CA 94720	1
	PRCF. I RUBIN SCHOOL OF ENGINEERING AND APPLIEC SCIENCE UNIVERSITY OF CALIFCRNIA LCS ANGELES, CALIFCRNIA 90024	1
*	PROF. I. R. SAVAGE DEPARTMENT OF STATISTICS YALE UNIVERSITY NEW HAVEN, CONNECTICUT C6520	1
	PROF. L. L., JR. SCHAFF DEFARTMENT OF ELECRICAL ENGINEERING COLORACO STATE UNIVERSITY FT. COLLINS , COLORADO 80521	ٌ ا
	PRCF. R. SERFLING DEPARTMENT CF STATISTICS FLORIDA STATE UNIVERSITY TALLAHASSEE FLORIDA 32306	1
	PROF. N. R. SCHUCANY DEPARTMENT OF STATISTICS SCUTHERN METHOLIST UNIVERSITY DALLAS , TEXAS 75222	1
	DEFT. CF STATISTICS STANFORD UNIVERSITY STANFORC CA 94305	. 1
	DEFT. CF FLECTRICAL ENGINEERING FOLYTECHNIC INSTITUTE OF NEW YORK BROOKLYN, NEW YORK 11201	1
	DEPT. CF CFERATIONS RESEARCH THE GEORGE WASHINGTON UNIVERSITY 707 22ND ST . N. W. WASHINGTON, C. C.	÷

L

,

4

÷

4

DR. A. L. SLAFKOSKY SCIENTIFIC ADVISOR COMMANCANT OF THE MARINE CORPS WASHINGTON , 1 C. C. 20380 MR. CHARLES S. SMITH MASD (IEL), FENT AGON WASHINGTON DC 1 20301 DR. D. E. SMITH DESMATICS INC. P.C. BCX 618 STATE COLLEGE PENNSYLVANIA J6801 1 FRCF. W. L. SMITH DEPARTMENT OF STATISTICS UNIVERSITY OF NORTH CARCLINA CHAPEL HILL * NOFTH CAROLINA 27514 1 FRCF. H SOLOMON DEFARTMENT OF STATISTICS STANFORD UNIVERSITY STANFORC , CALIFERNIA 94305 1 MR. GLENN F. STAHLY NATIONAL SECURITY AGENCY FORT MEADE MARYLANC 20755 1 MR. EAVIE A. SWICK AEVANCED FREJECIS GROUP CODE 8103 NAVAL FESEARCH LAB. MASHINGTEN DC 5 20375 WR. WENDELL G. SYKES ARTHUR D. LITTLE, INC. ACCRN PARK CAMBRICGE MA 1 C2140 FRUF. J. F. THOMPSON DEPARTMENT OF MATHEMATICAL SCIENCE RICE UNIVERSITY HOUSTON, TEXAS 77C01 1 .

.

.

ŧ

FRCF. L. A. THOMPSON DEPARTMENT OF STATISTICS UNIVERSITY OF MISSOURI COLUMBIA MISSOURI 65201	1
FRCF. F. A. TILLMAN DEPT. OF INCUSTRIAL ENGINEERING KANSAS STATE UNIVERSITY MANHATTAN KS 66506	1
FRCF J. W. TUKEY DEFAFIMENT OF STATISTICS PRINCETON UNIVERSITY PRINCETON , N. J. C854C	1
FROF. A F VEINOTT CEFARTMENT OF OPERATIONS RESEARCH STANFORD UNIVERSTITY STANFORD CALIFCRNIA 54305	1
CANIEL H. WAGNER STATION SCLARE ENE FAOLI, PENNSYLVANIA J9301	ì
PROF. GRACE WAIBA DEPT. OF STATISTICS UNIVERSITY OF WISCONSIN MADISON WI 53706	1
FRCF. K. T. WALLENIUS DEPARTMENT OF MATHEMATICAL SCIENCES CLEMSON UNIVERSITY CLEMSON. SOLTH CAROLINA 25631	1
PROF. G. S. LATSON CEPARTMENT OF STATISTICS FRINCETON, N. J. 08540	. 1
FREF. BERNARE WIEROW STANFERD ELECTRONIES LAB STANFERE UNIVERSITY STANFERD CA 543 05	1
FRCF. G. E. WHITEHOUSE DEPT. OF INDUSTRIAL ENGINEERING Lefigh UNIVERSITY Bethlehem FA	1

10

PROF. S. ZACKS DEFT. OF MATHEMATICS AND STATISTICS CASE WESTERN FESERVE UNIVERSITY CLEVELAND., CHIC 44106	1
PRCF. N. ZIA-FASSAN DEFARTMENT OF INDUSTRIAL AND SYSTEMS ENGINEERING ILLINCIS INSTITUTE OF TECHNOLOGY CFICAGO IL 60616	1
CR. DAVID BRILLINGER STATISTICS CEPT. UNIVERSITY CF CALIFORNIA BEFELEY. CALIFORNIA 54720	1
CR. CHIN CHEN CCMMONWELL TH ASSOCIATES, INC. 205 E. WASHINGTON JACKSON, MI 49201	1
FREFESSOR PETER CHEN ALFRED P. SLEAN SHEOOL OF MGNT MASS INSTITUTE OF TCHNCLOGY 50 MEMOFIAL DRIVE CAMBRIDGE MASSACHUSETTS 92139	1
DR. E. CINLAR O.R. DELLAR ACRTHWESTERN UNIVERSITY EVANSTON , ILLINDIS 60201	1
DR. GUY C CORYNEN L. 156 UNIV OF CALIFORNIA LAWRENCE LIVERMORE LABORATORY F. D. EOX EOE LIVEFMCRE, CALIFORNIA 94550	1
DR. MICHAEL A. CRANE CONTROL ANJLYSIS CORPORATION EOO WELCH PEAC FALD ALTE , CALIFORNIA 94304	1
CR. D. F. CALEY STATISTICS DEPT. (I A S) AUSTRALIAN NATIONAL UNIVERSITY CANBERRA, A. C. T. 2606 AUSTRALIA	1

1

1

1

ì

HSU DER-ANN SCH OF ENGF./APPLIED SCI. DEFT. OF CIVIL & GEOLOGICAL ENGR. THE ENGINEERING QUADRANGLE FRINCETON NEW JERSEY C8543 ASSISTANT CEAN CAV ELDREDGE SCHOOL OF EUSINESS ADMINISTRATION UNIVERSITY OF EVANSVILLE EVANSVILLE INDIANA 47702 CR. G. W. FISHMAN CIFRICULUM IN O. R. UNIVERSITY CF NORTH CAFCLINA PHILLIPS ANNEX , CHAPPEL HILL NORTH CARGLINA 27514 CR. A. V. FOLCEN DEPART HE NT OF PHYSIOLOGY THE UNIVERSITY LEEDS LS2 SJT ENGLANC 1 DR. ELGENE C. HOMER MANAGEMENT SCI. & ENGINEERING DEPT. C.W. POST COLLEGE CF LONG ISLAND UNIVERSITY GREENVALE, NEW YORK 1 11548 DR. D. VERE JONES DEPT. OF WATHEWATICS, VICTORIA UNIVERSITY OF WELLINGTON, P. O. BCX 196 WELLINGTON, NEW ZEALAND 1 ۰, J. M. LIITISCHAAGER SYSTEMS CIVISICA CCLLEGE CF ENGINEERING THE LNIVERSITY OF JOWA JOAA CITY, ICAA 52242 1 Y. W. CE KHAADSTENIET DE FAT FMENT OF EIULOGY FREE UNIVERSITY CE BOELELAAN 1087 AMSTERCAF-EUITENVELDER THE NETHERLANCS 1 JACK P. LANCOLT DEFENCE & CIVIL INST. OF ENVIR. MED 1133 SFEPART AVE. WEST PO BUX 2000 DOWNSVIEW ONTARIO, 1 CANADA

ł

*

The second second as a second second

- Contractor

MF. FRANK MCNOLTY 1142 PCME AVE. SUNNYVALE, CA 1 94087 CR. P. A. F. VCRAN STATISTICS CEPT. (I A S) AUSTRALIAN NATICNAL UNIVERSITY CANBERRF, A. C. T 2606 AUSTRALIA 1 G. W. MCRRISON DAK RIDGE NATICNAL LABORATORY CAK RICGE, TENNESSE 37830 1 MR. BRADLEY NICKEY LSCA FCREST SERVICE P. O. BOX 5007 FIVERSICE CALIFCRNIA 92507 1 CR. DAVID CAKES DEFT. OF STATISTICS HARVARC UNIVERSITY SCIENCE CENTER, 1 OXFORD ST. COMBRIDGE, MA. 02132 1 1 Professor Emanuel Parzen, Director Institute of Statistics. Teaxs A&M University College Station, TX 77843 . MR ART PETERSON DEFT. OF EIDSTATISTICS SC-32 UNOVERSITY OF WASHINGTON SEATTLE, WASHINGTON 1 Š8195 CAVID 6. RCBINSON CONTROL ANALYSIS CCRPOPATION 800 WELCF ROAC PALO ALTO CALIFORNIA 54204 1 C. H. SAUER(21-129) THOMAS J. WATSCN RESEARCH CENTER F.C. BCX 218 YCRKTUWN HEIGHTS, NEW YCRK 1 10598

.

1

1

1

1

1

1

.

WR. C. A. STEELE JR F.C. BCX 45 MAGNELIA, MA (1930 FRCF P. SULLO SCHOOL OF MANAGEMENT FENSSELAER POLYTECHNIC INSTITUTE TROY, N.Y. 12181 DR. FOREFT TRAPPL INSITIUT FUR ALLGEMEINE UNC VERGLFICHENDE PHYSIOLOGIE DER UNIVERSITAT WIEN SCHWARZSFANIEFSTRASSE 17 A-1090 WIEN OSTERREICH (ALSTRIA) PALL A. WILLIS PCLY TECHNICA 2824 W. GECRGE MASON FALLS CHURF , VA. 22042 TUNCEL M. YEGULALP KRLMB SCHCCL CF MINES 918 S.H. MLDD CCLUMBIA UNIVERSITY 520 H. 12CTF ST. NEW YROK; N.Y. 10027 SHIEGKI YOKOYAMA RESEARCH CENTER OF APPLIED INFORMATION SCIENCES TOHOKU UNIVERSITY KATAHIRA-CFO 2-1, SENDAI 980 JAPAN Dean of Research 1 **Code** 012 Naval Postgraduate School · Monterey, Ca. 93940 Library (Code 55) 1 Naval Postgraduate School Monterey, CA 93940

×.

NO. OF COPIESProfessor Peter Lewis50Code 55Lw50Naval Postgraduate School50Monterey, Ca. 9394950Mr. G. Shedler50K53-28250IBM Research Lab50Cottle Road5193R. J. Stampfel1Code 551Naval Postgraduate School1

•

Monterey, Ca. 93940