



Calhoun: The NPS Institutional Archive
DSpace Repository

Faculty and Researchers

Faculty and Researchers' Publications

2019

Modeling and Prediction of Arctic Climate Using the Regional Arctic System Model

Maslowski, Wieslaw; Osinski, Robert; Lee, Younjoo;
Kinney, Jaclyn Clement; Cassano, John; Seefeldt, Mark

Maslowski, Wieslaw, et al. "Modeling and Prediction of Arctic Climate Using the Regional Arctic System Model." Geophysical Research Abstracts. Vol. 21. 2019. <https://hdl.handle.net/10945/65922>

This publication is a work of the U.S. Government as defined in Title 17, United States Code, Section 101. Copyright protection is not available for this work in the United States.

Downloaded from NPS Archive: Calhoun



Calhoun is the Naval Postgraduate School's public access digital repository for research materials and institutional publications created by the NPS community. Calhoun is named for Professor of Mathematics Guy K. Calhoun, NPS's first appointed -- and published -- scholarly author.

Dudley Knox Library / Naval Postgraduate School
411 Dyer Road / 1 University Circle
Monterey, California USA 93943

<http://www.nps.edu/library>



Modeling and Prediction of Arctic Climate Using the Regional Arctic System Model

Wieslaw Maslowski (1), Robert Osinski (2), Younjoo Lee (1), Jaclyn Clement Kinney (1), John Cassano (3), and Mark Seefeldt (3)

(1) Naval Postgraduate School, Monterey, CA, USA (maslowsk@nps.edu), (2) Institute of Oceanology, Polish Academy of Sciences, Gdansk, Poland, (3) University of Colorado, Boulder, CO, USA

The Regional Arctic System Model (RASM) has been developed and used to investigate critical processes controlling the evolution of the Arctic climate system under a diminishing sea ice cover. RASM is a fully coupled limited-domain ice-ocean-atmosphere-land hydrology model. Its domain is pan-Arctic, with the atmosphere and land components configured on a 50-km or 25-km grid. The ocean and sea ice components are configured on rotated sphere meshes with four configuration options: 1/12o (~9.3km) or 1/48o (~2.4km) in the horizontal space and with 45 or 60 vertical layers. As a regional climate model, RASM requires boundary conditions along its lateral boundaries and in the upper atmosphere, which are derived either from global atmospheric reanalyses for simulations of the past to present or from Earth System models (ESMs) for climate projections. This allow comparison of RASM results with observations in place and time, which is a unique capability not available in global ESMs. Several examples of key physical processes and coupling between different model components will be presented, that improve the representation of the past and present Arctic climate system. The impact of such processes and feedbacks will be discussed with regard to improving model physics and reducing biases in the representation of its initial state for prediction of Arctic climate change at time scales from synoptic to decadal.

Copyright of Geophysical Research Abstracts is the property of Copernicus Gesellschaft mbH and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.