



Calhoun: The NPS Institutional Archive
DSpace Repository

Faculty and Researchers

Faculty and Researchers' Publications

2019-05

Process-resolving Regional Arctic System Model for Advanced Modeling and Prediction of Arctic Climate System

Maslowski, Wieslaw; Osinski, Robert; Lee, Younjoo; Kinney, Jaclyn Clement; Cassano, J.J.; Seefeldt, Mark W.; Craig, Anthony P.; Nijssen, Bart; Gergel, Diana R.

AMS

Maslowski, Wieslaw, et al. "Process-resolving Regional Arctic System Model for Advanced Modeling and Prediction of Arctic Climate System." 15th Conference on Polar Meteorology and Oceanography. AMS, 2019.

<http://hdl.handle.net/10945/66156>

This publication is a work of the U.S. Government as defined in Title 17. United



Downloaded from NPS Archive: Calhoun

Calhoun is the Naval Postgraduate School's public access digital repository for research materials and institutional publications created by the NPS community. Calhoun is named for Professor of Mathematics Guy K. Calhoun, NPS's first appointed -- and published -- scholarly author.

Dudley Knox Library / Naval Postgraduate School
411 Dyer Road / 1 University Circle
Monterey, California USA 93943

<http://www.nps.edu/library>



7A.2: Process-resolving Regional Arctic System Model for Advanced Modeling and Prediction of Arctic Climate System

Tuesday, May 21, 2019

01:45 PM - 02:00 PM

📍 *Williams Village Conference Center - Room C*

The Regional Arctic System Model (RASM) is a fully coupled limited-domain ice-ocean-atmosphere-land hydrology model. Its domain is pan-Arctic, with the atmosphere and land components configured on a 50-km or 25-km grid. The ocean and sea ice components are configured on rotated sphere meshes with four configuration options: $1/12^\circ$ (~9.3km) or $1/48^\circ$ (~2.4km) in the horizontal space and with 45 or 60 vertical layers. As a regional climate model, RASM requires boundary conditions along its lateral boundaries and in the upper atmosphere, which are derived either from global atmospheric reanalyses for simulations of the past to present or from Earth System models (ESMs) for climate projections. In the former case, this allow comparison of RASM results with observations in place and time, which is a unique capability not available in global ESMs. RASM has been developed and used to investigate critical processes controlling the evolution of the Arctic climate system under a diminishing sea ice cover. Several examples of key physical processes and coupling between different model components will be presented, that improve the representation of the past and present Arctic climate system. The impact of such processes and feedbacks will be discussed with regard to improving model physics and reducing biases in the representation of its initial state for prediction of Arctic climate at time scales from synoptic to intra-annual.

Authors

Wieslaw Maslowski

*Naval Postgraduate School
Monterey, CA, USA*

Robert Osinski

*Institute of Oceanology, Polish
Academy of Sciences
Sopot, Poland*

Younjoo Lee

*Naval Postgraduate School
Monterey, CA, USA*

Jaclyn Clement Kinney

*Naval Postgraduate School
Monterey, CA, USA*

J.J. Cassano

*CIRES/Univ. of Colorado
Boulder, CO, USA*

Mark W. Seefeldt

*CIRES/Univ. of Colorado
Boulder, CO, USA*

Anthony P. Craig

*NPS Contractor
Monterey, CA, USA*

Bart Nijssen

*Univ. of Washington
Seattle, WA, USA*

Diana R. Gergel

*Univ. of Washington
Seattle, WA, USA*



 **Recorded Presentation**
Recording.

[Find Similar](#)

View Related Events

Day: Tuesday, May 21, 2019