



**Calhoun: The NPS Institutional Archive**  
**DSpace Repository**

---

NPS Scholarship

Publications

---

2014-06

## Sopite Syndrome: A Revised Definition

Matsangas, Panagiotis; McCauley, Michael E.

---

Matsangas, P. McCauley, M.E. Sopite syndrome: a revised definition. *Aviation, Space, and Environmental Medicine*, Volume 85, No. 6, June 2014 pp. 672-3.  
<https://hdl.handle.net/10945/45394>

---

This publication is a work of the U.S. Government as defined in Title 17, United States Code, Section 101. Copyright protection is not available for this work in the United States.

*Downloaded from NPS Archive: Calhoun*



Calhoun is the Naval Postgraduate School's public access digital repository for research materials and institutional publications created by the NPS community. Calhoun is named for Professor of Mathematics Guy K. Calhoun, NPS's first appointed -- and published -- scholarly author.

**Dudley Knox Library / Naval Postgraduate School**  
**411 Dyer Road / 1 University Circle**  
**Monterey, California USA 93943**

<http://www.nps.edu/library>

# Sopite Syndrome: A Revised Definition

PANAGIOTIS MATSANGAS AND MICHAEL E. MCCAULEY

MATSANGAS P, MCCAULEY ME. *Sopite syndrome: a revised definition. Aviat Space Environ Med.* 2014; 85:672-3.

In 1976, Graybiel and Knepton proposed the term "sopite syndrome" to describe a symptom complex centering on drowsiness and lethargy related to motion sickness. However, existing descriptions and definitions of sopite syndrome have limitations in fully conveying the appropriate information to the reader. Our objective is to propose a revised definition providing a more adequate conceptual framework for research. The proposed definition of sopite syndrome addresses the nonspecificity of soporific symptoms, the health state of the individuals, and the existence of a motion stimulus.

**Keywords:** motion effects, motion sickness, drowsiness.

IT HAS LONG BEEN KNOWN that symptoms like drowsiness, apathy, lassitude, lethargy, and lack of interest in the task or the environment occur in nauseogenic motion conditions (5,12). Research conducted in a slow rotating room (SRR) reported depression and sleepiness (3). Reason and Brand (12) reported an unpublished study by Reason and Graybiel involving a 3-d exposure to angular accelerations where drowsiness was persistent and overwhelming.

Graybiel and Knepton (4) identified and described "sopite syndrome," a symptom-complex centering on drowsiness and lethargy related to motion sickness. Sopite syndrome is a response to motion that includes symptoms such as drowsiness, lethargy, apathy, decreased ability to concentrate, daydreaming, melancholy, yawning, disinterest and disinclination to work, lack of participation in group activities, mood changes, irritability, sleep disturbances, frequent daytime napping, mild depression, and a desire to be left alone (4,9). This description was followed by ISO 5805, in which sopite syndrome was defined as "inordinate sleepiness, lassitude or drowsy inattention induced by vibration, low-frequency oscillatory motion (e.g., ship motion) or general travel stress..." (6). Depending on the stimulus, sopite syndrome may be the only reported manifestation of motion sickness (4,11). Soporific symptoms have been shown to appear before nausea and to remain after the cessation of a nauseogenic motion stimulus (9).

Sopite syndrome is important operationally for a number of reasons: a) a person with sopite syndrome may have degraded performance, but may not be identified as motion sick; b) drugs for nausea may not improve degraded performance due to sopite effects; and c) medication side effects might exacerbate existing drowsiness due to sopite syndrome (1,4,9). Soporific effects are operationally important because they are common and frequent. Research has shown that drowsiness is among the most frequent symptoms associated with motion sickness (2). Lastly, it has been suggested that sopite syndrome can lead to inefficiency and being accident

prone and that it could have profound effects in transport environments where, for other reasons, sleep disturbances exist (4,9).

Graybiel and Knepton's (4) work describes evidence from earlier research, identifies the major attributes of the syndrome, and discusses why there is a need to define a symptom-complex centering around drowsiness. However, their work did not provide a structured definition of sopite syndrome. Existing descriptions/definitions have limitations in fully conveying the appropriate information to the reader. We identified three issues of concern: nonspecificity of soporific symptoms; the health state of the individuals; and the existence of a motion stimulus.

Symptoms and signs of motion sickness and sopite syndrome are nonspecific (4,7,8), i.e., they also can be observed in the absence of a nauseogenic stimulus, because of stress, illness or other reasons.

Hill (5) discussed the association between seasickness and drowsiness, apathy, and mental lethargy, emphasizing that these symptoms occur in the absence of actual somnolence. Similarly, Lawson and Mead (9) indicated that sopite syndrome is distinct from the state of fatigue. The ISO 5805 definition addressed this issue by noting the "inordinate" attribute of sleepiness and lassitude when referring to soporific symptoms. Underlying pathological conditions may lead to many of the symptoms associated with motion sickness, for example nausea because of stress, or lethargy because of depression, etc. The investigation of motion sickness with healthy individuals partially alleviates the problem of nonspecificity, to the extent that this relates to pathognomonic reasons. Therefore, nonspecificity of symptoms is a critical issue when discussing sopite syndrome and should be emphasized in the corresponding definition.

Furthermore, the existence of a motion stimulus is a causal factor when discussing sopite syndrome (4,9). Soporific symptomatology is evident both in real and apparent motion settings (7,9,10). Yet the definition

From the Department of Operations Research, Naval Postgraduate School, Monterey, CA.

This manuscript was received for review in October 2013. It was accepted for publication in February 2014.

Address correspondence and reprint requests to: Panagiotis Matsangas, 1411 Cunningham Road, Naval Postgraduate School, Monterey, CA 93943-5219; pmatsang@nps.edu.

Reprint & Copyright © by the Aerospace Medical Association, Alexandria, VA.

DOI: 10.3357/ASEM.3891.2014

provided in ISO 5805 does not clarify that sopite syndrome can be observed even in optokinetic stimulation environments (6). Furthermore, the ISO 5805 definition associates sopite syndrome with "... general travel stress." However, the effect of general stress on the development of sopite syndrome is not confirmed by existing research (4,9). Existing research indicates that sopite syndrome often may be confounded by other forms of fatigue coexisting in operational settings (e.g., airplanes, cars, trains) because of stress or other reasons (9). Therefore, sopite syndrome should not be confused with other stressors.

Our objective is to propose the following revised definition of sopite syndrome which addresses the aforementioned shortfalls:

Sopite syndrome is a symptom complex that develops as a result of exposure to real or apparent motion and is characterized by excessive drowsiness, lassitude, lethargy, mild depression, and reduced ability to focus on an assigned task. Sopite syndrome is most clearly distinguished in a healthy individual free from pathological conditions that engender similar symptoms and not suffering from sleep deprivation, mental or physical fatigue, or increased levels of physical activity.

In summary, existing descriptions and definitions of sopite syndrome raise a number of concerns. The definition given above addresses the limitations of earlier approaches and provides an adequate conceptual framework for research.

#### ACKNOWLEDGMENTS

*Authors and affiliations:* Panagiotis Matsangas, Ph.D. and Michael McCauley, Ph.D., Department of Operations Research, Naval Postgraduate School, Monterey, CA.

#### REFERENCES

1. Buckley JC, Jr. Space physiology. New York, NY: Oxford University Press; 2006:187-206.
2. Cowings PS, Toscano WB, DeRoshia C, Tauson RA. Effects of Command and Control Vehicle (C2V) operational environment on soldier health and performance. *Hum Perf Extrem Environ* 2001; 5:66-91.
3. Graybiel A, Clark B, Zariello JJ. Observations on human subjects living in a "slow rotation room" for periods of two days. *Arch Neurol* 1960; 3:55-73.
4. Graybiel A, Knepton J. Sopite syndrome: a sometimes sole manifestation of motion sickness. *Aviat Space Environ Med* 1976; 47:873-82.
5. Hill J. The care of the sea-sick. *BMJ* 1936; 2:802-7.
6. International Organization for Standardization. Mechanical vibration and shock—Human exposure – Vocabulary (ISO 5805:1997): International Organization for Standardization; 1997.
7. Kiniorski ET, Weider SK, Finley JR, Fitzgerald EM, Howard JC, Di Nardo PA. Sopite symptoms in the optokinetic drum. *Aviat Space Environ Med* 2004; 75:872-5.
8. Lang IM, Sarna SK, Shaker R. Gastrointestinal motor and myoelectric correlates of motion sickness. *Am J Physiol* 1999; 277:G642-52.
9. Lawson BD, Mead AM. The sopite syndrome revisited: drowsiness and mood changes during real or apparent motion. *Acta Astronaut* 1998; 43:181-92.
10. Matsangas P. The effect of mild motion sickness and sopite syndrome in cognitive multitasking performance [Dissertation]. Monterey, CA: Naval Postgraduate School; 2013.
11. Mead AM, Lawson BD. Sopite syndrome case report I: Motion-induced drowsiness and mood changes in an individual with no other motion sickness symptoms – a case of "pure" sopite syndrome? [abstract]. *Aviat Space Environ Med* 1997; 68:648.
12. Reason JT, Brand JJ. Motion sickness. Oxford, England: Academic Press; 1975.

