



Calhoun: The NPS Institutional Archive
DSpace Repository

NPS Scholarship

Publications

1995-07-03

Precision measurement of the deuteron spin structure function gd_1

Abe, K.; Akagi, T.; Anthony, P.L.; Antonov, R.; Arnold, R.G.; Averett, T.; Band, H.R.; Bauer, J.M.; Borel, H.; Bosted, P.E....

American Physical Society

Physical Review Letters, v. 75, no.1 (July 3, 1995), pp. 25-28
<https://hdl.handle.net/10945/48044>

This publication is a work of the U.S. Government as defined in Title 17, United States Code, Section 101. Copyright protection is not available for this work in the United States.

Downloaded from NPS Archive: Calhoun



Calhoun is the Naval Postgraduate School's public access digital repository for research materials and institutional publications created by the NPS community. Calhoun is named for Professor of Mathematics Guy K. Calhoun, NPS's first appointed -- and published -- scholarly author.

Dudley Knox Library / Naval Postgraduate School
411 Dyer Road / 1 University Circle
Monterey, California USA 93943

<http://www.nps.edu/library>

Precision Measurement of the Deuteron Spin Structure Function g_1^d

K. Abe,¹⁵ T. Akagi,^{12,15} P. L. Anthony,¹² R. Antonov,¹¹ R. G. Arnold,¹ T. Averett,¹⁶ H. R. Band,¹⁷ J. M. Bauer,⁷ H. Borel,⁵ P. E. Bosted,¹ V. Breton,³ J. Button-Shafer,⁷ J. P. Chen,⁴ T. E. Chupp,⁸ J. Clendenin,¹² C. Comptour,³ K. P. Coulter,⁸ G. Court,^{12,*} D. Crabb,¹⁶ M. Daoudi,¹² D. Day,¹⁶ F. S. Dietrich,⁶ J. Dunne,¹ H. Dutz,^{12,†} R. Erbacher,^{12,13} J. Fellbaum,¹ A. Feltham,² H. Fonvieille,³ E. Frlez,¹⁶ D. Garvey,⁹ R. Gearhart,¹² J. Gomez,⁴ P. Grenier,⁵ K. A. Griffioen,^{11,‡} S. Hoibraten,^{16,§} E. W. Hughes,¹² C. Hyde-Wright,¹⁰ J. R. Johnson,¹⁷ D. Kawall,¹³ A. Klein,¹⁰ S. E. Kuhn,¹⁰ M. Kuriki,¹⁵ R. Lindgren,¹⁶ T. Liu,¹⁶ R. M. Lombard-Nelsen,⁵ J. Marroncle,⁵ T. Maruyama,¹² X. K. Maruyama,⁹ J. McCarthy,¹⁶ W. Meyer,^{12,†} Z.-E. Meziani,^{13,14} R. Minehart,¹⁶ J. Mitchell,⁴ J. Morgenstern,⁵ G. G. Petratos,^{12,||} R. Pitthan,¹² D. Pocanic,¹⁶ C. Prescott,¹² R. Prepost,¹⁷ P. Raines,¹¹ B. Raue,¹⁰ D. Reyna,¹ A. Rijllart,^{12,¶} Y. Roblin,³ L. S. Rochester,¹² S. E. Rock,¹ O. A. Rondon,¹⁶ I. Sick,² L. C. Smith,¹⁶ T. B. Smith,⁸ M. Spengos,¹ F. Staley,⁵ P. Steiner,² S. St. Lorant,¹² L. M. Stuart,¹² F. Suekane,¹⁵ Z. M. Szalata,¹ H. Tang,¹² Y. Terrien,⁵ T. Usher,¹² D. Walz,¹² J. L. White,¹ K. Witte,¹² C. C. Young,¹² B. Youngman,¹² H. Yuta,¹⁵ G. Zapalac,¹⁷ B. Zihlmann,² and D. Zimmermann¹⁶

(E143 Collaboration)

¹The American University, Washington, D.C. 20016

²Institut für Physik der Universität Basel, CH 4056 Basel, Switzerland

³Laboratoire de Physique Corpusculaire, IN2P3/CNRS, University Blaise Pascal, F-63170 Aubiere Cedex, France

⁴Continuous Electron Beam Accelerator Facility, Newport News, Virginia 23606

⁵DAPNIA-Service de Physique Nucleaire Centre d'Etudes de Saclay, 91191 Gif-sur-Yvette, France

⁶Lawrence Livermore National Laboratory, Livermore, California 94550

⁷University of Massachusetts, Amherst, Massachusetts 01003

⁸University of Michigan, Ann Arbor, Michigan 48109

⁹Naval Postgraduate School, Monterey, California 93943

¹⁰Old Dominion University, Norfolk, Virginia 23529

¹¹University of Pennsylvania, Philadelphia, Pennsylvania 19104

¹²Stanford Linear Accelerator Center, Stanford, California 94309

¹³Stanford University, Stanford, California 94305

¹⁴Temple University, Philadelphia, Pennsylvania 19122

¹⁵Tohoku University, Sendai 980, Japan

¹⁶University of Virginia, Charlottesville, Virginia 22901

¹⁷University of Wisconsin, Madison, Wisconsin 53706

(Received 8 March 1995)

We report on a high-statistics measurement of the deuteron spin structure function g_1^d at a beam energy of 29 GeV in the kinematic range $0.029 < x < 0.8$ and $1 < Q^2 < 10$ (GeV/c)². The integral $\Gamma_1^d = \int_0^1 g_1^d dx$ evaluated at fixed $Q^2 = 3$ (GeV/c)² gives $0.042 \pm 0.003(\text{stat}) \pm 0.004(\text{syst})$. Combining this result with our earlier measurement of g_1^p , we find $\Gamma_1^p - \Gamma_1^n = 0.163 \pm 0.010(\text{stat}) \pm 0.016(\text{syst})$, which agrees with the prediction of the Bjorken sum rule with $O(\alpha_s^3)$ corrections, $\Gamma_1^p - \Gamma_1^n = 0.171 \pm 0.008$. We find the quark contribution to the proton helicity to be $\Delta q = 0.30 \pm 0.06$.

PACS numbers: 25.30.Fj, 11.55.Hx, 13.60.Hb, 13.88.+e

The longitudinal and transverse spin-dependent structure functions $g_1(x, Q^2)$ and $g_2(x, Q^2)$ for polarized deep-inelastic lepton-nucleon scattering provide information on the spin structure of the proton and neutron. A fundamental QCD sum rule, originally derived from current algebra by Bjorken [1], predicts the difference $\Gamma_1^p - \Gamma_1^n = \frac{1}{6}(g_A/g_V)$ at infinite Q^2 where $\Gamma_1^{p(n)} = \int_0^1 g_1^{p(n)}(x, Q^2) dx$ for the proton (neutron), and g_A and g_V are the axial-vector and vector coupling constants in neutron β decay. QCD corrections up to third order in α_s have been computed [2], thus making a test of the Bjorken sum rule possible at finite Q^2 . Measurements of Γ_1^p [3,4], Γ_1^n

from ³He [5], as well as results from deuterium [6] targets, are in agreement with this prediction within experimental uncertainties. Separate sum rules for Γ_1^p and Γ_1^n were derived by Ellis and Jaffe [7] under the assumptions of SU(3) flavor symmetry and an unpolarized strange sea. Higher order QCD corrections have been calculated [8,9].

The polarized spin structure function g_1 is related to the virtual photon asymmetries A_1 and A_2 :

$$g_1 = \frac{F_1}{(1 + \gamma^2)}(A_1 + \gamma A_2), \quad (1)$$

where F_1 is the unpolarized structure function, $\gamma^2 = Q^2/\nu^2$, $\nu = E - E'$, E and E' are incident and scattered electron energies, respectively, and A_1 and A_2 are virtual photon cross section asymmetries [10]. These asymmetries are related by kinematic factors to the experimentally measured electron asymmetries A_{\parallel} and A_{\perp} . The longitudinal asymmetry A_{\parallel} is the cross section asymmetry between negative- and positive-helicity electron beams when the target nucleon is polarized parallel to the beam direction. The transverse asymmetry A_{\perp} is the asymmetry when the target nucleon is polarized transverse to the beam direction. We use the relationships $A_1 = (A_{\parallel}/D - \eta A_{\perp}/d)/(1 + \eta\zeta)$ and $A_2 = (\zeta A_{\parallel}/D + A_{\perp}d)/(1 + \eta\zeta)$, where $\eta = \epsilon\sqrt{Q^2/E^2}/(1 - \epsilon E'/E)$, $\zeta = \eta(1 + \epsilon)/2\epsilon$, $D = (1 - \epsilon E'/E)/(1 + \epsilon R)$ is the depolarization factor, $d = D\sqrt{2\epsilon/(1 + \epsilon)}$, $\epsilon^{-1} = 1 + 2[1 + (\nu^2/Q^2)]\tan^2(\theta/2)$, $R = \sigma_L/\sigma_T$, and θ is the electron scattering angle. Thus, the quantity g_1 can be written as $g_1 = (F_1/D')[A_{\parallel} + \tan(\theta/2)A_{\perp}]$, where $D' = (1 - \epsilon)(2 - y)/y(1 + \epsilon R)$, and $y = \nu/E$.

Experiment E143 used the SLAC polarized electron beam with energies of 9.7, 16.2, and 29.1 GeV incident on polarized proton and deuteron targets in End Station A to measure g_1^p , g_2^p , g_1^d , and g_2^d in the range $1 < Q^2 < 10$ (GeV/c)² and $0.029 < x < 0.8$. This Letter reports on our analysis of the 29.1 GeV data, which yielded g_1^d results with considerably smaller statistical uncertainties than previous measurements [6]. We adopt the convention that g_1^d refers to the average structure function of the nucleon in the deuteron: $g_1^d \approx \frac{1}{2}(g_1^p + g_1^n)$.

The helicity of the longitudinally polarized electron beam [11] was selected randomly on a pulse-to-pulse basis to minimize instrumental asymmetries, which were found to be negligible. The beam polarization P_b was measured daily with a Møller polarimeter, and was found to vary with the cathode quantum efficiency from 0.83 to 0.86. An overall uncertainty on P_b of ± 0.02 was achieved [4].

The target [12] was a 3-cm-long 2.5-cm-diameter cylinder filled with granules of deuterated ammonia, ¹⁵ND₃, of greater than 98% isotopic purity. It was polarized by the technique of dynamic nuclear polarization in a 4.8-T magnetic field. An average in-beam polarization P_t of 25% was measured with an NMR technique, and a maximum of greater than 40% was achieved. The NMR signal was calibrated by measuring the thermal-equilibrium polarization near 1.6 K. An overall relative uncertainty of 4% on P_t was achieved.

Scattered electrons were detected in two independent spectrometers [13] at angles of 4.5° and 7° with respect to the incident beam. Electrons were identified in each spectrometer by use of two threshold gas Čerenkov counters and a 200-element shower-counter array of lead glass blocks 24 radiation lengths thick. Particle momenta and scattered angles were measured with seven planes of scintillator hodoscopes.

The experimental longitudinal and transverse asymmetries A_{\parallel} and A_{\perp} were determined from

$$A_{\parallel} \text{ (or } A_{\perp}) = C_1 \left(\frac{N_L - N_R}{N_L + N_R} \frac{1}{fP_bP_t} - C_2 \right) + A_{rc}, \quad (2)$$

where N_L and N_R are the corrected numbers of scattered electrons per incident charge for negative and positive beam helicity, respectively. Charge-symmetric backgrounds measured by reversing the spectrometer polarity have no measurable asymmetry. They led to rate corrections of 10% at the lowest x bin, decreasing rapidly with x . The rates were also corrected for dead time effects.

The correction factors C_1 and C_2 account for the polarizations of ¹⁵N, unsubstituted ¹⁴N, and residual protons in the target. The factor C_2 ranges from 3% to 5% of the measured proton asymmetry, and C_1 is typically 1.016. These factors were determined from measured nitrogen polarizations and a shell-model calculation to determine the contribution of the unpaired p -shell proton.

The dilution factor f represents the fraction of measured events expected to originate from polarizable deuterons in the target. It was calculated from the composition of the target, which contained about 23% deuterons, 56% ¹⁵N, 10% ⁴He, 6% Al, 4% Cu, and 1% Ti by weight. The dilution factor varied from 0.22 at low x to 0.25 at high x . The relative systematic error in f was determined from uncertainties in the target composition and cross section ratios, and ranged from 2.2% to 2.6%.

The radiative correction A_{rc} includes both internal [14] and external [15] contributions, and typically changed A_{\parallel} by 10% of its value. Systematic errors on A_{rc} were estimated based on uncertainties in the input models and correspond to relative errors on A_{\parallel} of typically 7% for $x > 0.1$, increasing to 100% at $x = 0.03$.

Data from the two spectrometers, which differ by about a factor of 2 in average Q^2 , are consistent with A_1^d , A_2^d , and g_1^d/F_1^d being independent of Q^2 in the overlap region $0.07 < x < 0.55$, and therefore have been averaged together. The values of A_1^d from this experiment at $E = 29.1$ GeV shown in Fig. 1 [16] are consistent with the higher Q^2 results from the SMC Collaboration [6].

Values of g_1^d at the average $Q^2 = 3$ (GeV/c)² of this experiment are shown in Fig. 2(a). The evaluation of g_1^d at constant Q^2 is model dependent. We made the assumption that g_1^d/F_1^d is independent of Q^2 . For $F_1^d/(1 + \gamma^2) = F_2^d/2x(1 + R)$ we used the NMC fit [17] to F_2^d and the SLAC fit to R [18]. Using the SLAC fit to F_2^d [19] gives similar results. The systematic error on F_1^d is typically 2.5%, increasing to 5% at the lowest x bin and 15% at the highest x bin. The integral of g_1^d at $Q^2 = 3$ (GeV/c)² and over the measured x range is $\int_{0.029}^{0.8} g_1^d(x) dx = 0.040 \pm 0.003(\text{stat}) \pm 0.004(\text{stat})$. The integral is decreased by 0.002 if we make an alternate assumption that A_1^d and A_2^d are independent of Q^2 .




FIG. 1. The virtual photon asymmetry A_1^d from this experiment. The systematic errors are indicated by the shaded band. The average Q^2 varies from 1.3 $(\text{GeV}/c)^2$ at low x to 9 $(\text{GeV}/c)^2$ at high x . Data from the SMC Collaboration [6] are also shown.

Assuming g_1^d varies as $(1-x)^3$ at high x [20], the extrapolation for $x > 0.8$ yields $\int_{0.8}^1 g_1^d(x) dx = 0.000 \pm 0.001$. To make the extrapolation to $x = 0$, we make a model-dependent assumption that the data are described by the Regge-motivated form [21] $g_1^d(x) = Cx^{-\alpha}$, where α is allowed to be in the range -0.5 to 0 [22]. A fit to the data of this experiment in the range $x < x_{\text{max}} = 0.1$ gives $\int_0^{0.029} g_1^d(x) dx = 0.001 \pm 0.001$. The uncertainty includes a statistical component, the uncertainty in α , and the effect of varying the fitting range from $x_{\text{max}} = 0.05$ to $x_{\text{max}} = 0.12$. Including the SMC data [6] in the fit does not change the results. An alternate form [23], $g_1^d(x) = C' \ln(x)$, which provides good fits to low x data




FIG. 2. Values of xg_1 from this experiment (E143) as a function of x for (a) the deuteron and (b) the neutron. The errors are statistical only. Systematic errors are indicated by the shaded bands. Also shown are the neutron results from SLAC E142 [5].

on F_2 from HERA, gives similar results. Thus, we obtain the total integral $\Gamma_1^d(E143) = 0.042 \pm 0.003(\text{stat}) \pm 0.004(\text{syst})$, to be compared with the results of SMC at $Q^2 = 4.7 (\text{GeV}/c)^2$: $\Gamma_1^d(\text{SMC}) = 0.023 \pm 0.020(\text{stat}) \pm 0.015(\text{syst})$. The contributions to systematic uncertainties are given in Table I.

The Ellis-Jaffe sum rule for the deuteron is related to the sum rules for the proton and the neutron by $\Gamma_1^d = \frac{1}{2}(\Gamma_1^p + \Gamma_1^n)(1 - 1.5\omega_D)$, where ω_D is the probability that the deuteron is in a D state. We use $\omega_D = 0.05 \pm 0.01$ [24] given by N - N potential calculations. No other nuclear contributions to ω_D are included. The sum rule predicts $\Gamma_1^d(\text{EJ}) = 0.069 \pm 0.004$, where we have used $F + D = g_A/g_V = 1.2573 \pm 0.0028$ and $F/D = 0.575 \pm 0.016$ [25], and $\alpha_s = 0.35 \pm 0.05$ [26] for QCD corrections [8] to $Q^2 = 3 (\text{GeV}/c)^2$. Our measurement of Γ_1^d provides a precise test of the Ellis-Jaffe sum rule, and shows a disagreement of more than 3 standard deviations.

The spin structure function integral can be used in the quark parton model to extract the helicity contributions to the proton of each type of quark and antiquark. Measurements using the deuteron are expected to be more sensitive than those using either the proton or the neutron [27]. We find the total contribution from all quarks to be $\Delta q = 0.30 \pm 0.06$, and the contribution from strange quarks and antiquarks to be $\Delta s = -0.09 \pm 0.02$. They are the most precise determinations to date and are consistent with earlier results [28].

The neutron spin structure function can be extracted using the relation $g_1^n(x) = 2g_1^d(x)/(1 - 1.5\omega_D) - g_1^p(x)$. The results obtained using our earlier measurements [4] of $g_1^p(x)$ are compared in Fig. 2(b) with the results obtained by E142 [5] at $Q^2 = 2 (\text{GeV}/c)^2$ using a ^3He target. Using the same extrapolation procedure, we find $\Gamma_1^n(\text{E143}) = -0.037 \pm 0.008(\text{stat}) \pm 0.011(\text{syst})$, compared with $\Gamma_1^n(\text{E142}) = -0.022 \pm 0.007(\text{stat}) \pm 0.009(\text{syst})$. The correlations between proton and deuteron measurements were accounted for when determining the systematic uncertainty contributions in Table I.

The Bjorken sum rule prediction of $\Gamma_1^p - \Gamma_1^n = 0.171 \pm 0.008$ at $3 (\text{GeV}/c)^2$ can be tested by combining

TABLE I. Contributions to the systematic uncertainties of Γ_1^d , Γ_1^n , and $\Gamma_1^p - \Gamma_1^n$.

Uncertainty	$\delta\Gamma_1^d$	$\delta\Gamma_1^n$	$\delta(\Gamma_1^p - \Gamma_1^n)$
P_b	0.001	0.001	0.004
P_t	0.002	0.005	0.007
f	0.002	0.006	0.008
A_{rc}	0.002	0.006	0.007
F_2 and R	0.001	0.002	0.005
Extrapolation	0.001	0.004	0.006
Total	0.004	0.011	0.016

proton and deuteron results: $g_1^p(x) - g_1^n(x) = 2g_1^p(x) - 2g_1^d(x)/(1 - 1.5\omega_D)$. Following the same extrapolation procedure as in the case of the deuteron, we find $\Gamma_1^p(E143) - \Gamma_1^n(E143) = 0.163 \pm 0.010(\text{stat}) \pm 0.016(\text{syst})$, consistent with the prediction. Contributions to systematic uncertainties are given in Table I. This result is also consistent with that obtained by combining the Γ_1^p result from this experiment [4] and the Γ_1^n result from E142 [5]: $\Gamma_1^p(E143) - \Gamma_1^n(E142) = 0.149 \pm 0.014$.

In conclusion, we have performed a high-statistics measurement of the deuteron spin structure function g_1^d , and find that it disagrees with the Ellis-Jaffe sum rule by more than 3 standard deviations. When combined with our earlier results on g_1^p , we find the difference $\Gamma_1^p - \Gamma_1^n$ is in agreement with the fundamental Bjorken sum rule.

We wish to acknowledge the tremendous effort made by the SLAC staff in making this experiment successful; H. Wiedemann and SSRL, and the staffs of the Bates Linear Accelerator, CEBAF, and the Saskatoon accelerator for providing beams to irradiate the ammonia targets; and N. Shumeiko for help with radiative corrections. This work was supported by Department of Energy Contract No. DE-AC05-84ER40150 (CEBAF), No. W-2705-Eng-48 (LLNL), No. DE-AC03-76SF00515 (SLAC), No. DE-FG03-88ER 40439 (Stanford), No. DE-FG05-94ER40859 (ODU), No. DE-FG05-88ER40390 and No. DE-FG05-86ER40261 (Virginia), and No. DE-AC02-76ER00881 (Wisconsin); by National Science Foundation Grants No. 9114958 (American), No. 9307710 (Massachusetts), No. 9217979 (Michigan), No. 9104975 (ODU), and No. 9118137 (U. Penn.); by the Schweizersche Nationalfonds (Basel); by the Commonwealth of Virginia (Virginia); by the Centre National de la Recherche Scientifique (LPC Clermont-Ferrand); by the Commissariat a l'Energie Atomique (CE Saclay); and by the Ministry of Science, Culture and Education of Japan (Tohoku).

*Permanent address: Oliver Lodge Laboratory, University of Liverpool, Liverpool, U.K.

†Permanent address: University of Bonn, Bonn, D-53313 Germany.

‡Present address: College of William and Mary, Williamsburg, VA 23187.

§Permanent address: FFIYM, P.O. Box 25, N-2007 Kjeller, Norway.

||Present address: Kent State University, Kent, OH 44242.

¶Permanent address: CERN, 1211 Geneva 23, Switzerland.

- [1] J. D. Bjorken, Phys. Rev. **148**, 1467 (1966); Phys. Rev. D **1**, 1376 (1970).
 [2] S. A. Larin and J. A. M. Vermaseren, Phys. Lett. B **259**, 345 (1991), and references therein.
 [3] SLAC E80 Collaboration, M. J. Alguard *et al.*, Phys. Rev. Lett. **37**, 1261 (1976); **41**, 70 (1978); SLAC E130

- Collaboration, G. Baum *et al.*, *ibid.* **51**, 1135 (1983); EMC Collaboration, J. Ashman *et al.*, Phys. Lett. B **206**, 364 (1988); Nucl. Phys. **B328**, 1 (1989); SMC Collaboration, D. Adams *et al.*, Phys. Lett. B **329**, 399 (1994).
 [4] SLAC E143 Collaboration, K. Abe *et al.*, Report No. SLAC-PUB-6508, 1994 (unpublished); Phys. Rev. Lett. **74**, 346 (1995).
 [5] SLAC E142 Collaboration, P. L. Anthony *et al.*, Phys. Rev. Lett. **71**, 959 (1993).
 [6] SMC Collaboration, B. Adeva *et al.*, Phys. Lett. B **302**, 533 (1993).
 [7] J. Ellis and R. L. Jaffe, Phys. Rev. D **9**, 1444 (1974); **10**, 1669 (1974).
 [8] S. Larin, Phys. Lett. B **334**, 192 (1994).
 [9] A. L. Kataev, Report No. CERN-TH-7427/94, 1994 (unpublished).
 [10] R. D. Roberts, *The Structure of the Proton* (Cambridge University Press, Cambridge, 1990).
 [11] T. Maruyama, E. Garwin, R. Prepost, and G. Zapalac, Phys. Rev. B **46**, 4261 (1992); D. Schultz *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **340**, 127 (1994).
 [12] D. G. Crabb and D. Day, in *Proceedings of the 7th Workshop on Polarized Target Materials and Techniques, Bad Honnef, Germany, 1994* [Nucl. Instrum. Methods Phys. Res., Sect. A **9** (1995)].
 [13] G. G. Petratos *et al.*, Report No. SLAC-PUB-5678, 1991 (unpublished); H. Borel *et al.*, in IEEE Conference, Norfolk, VA, 1994 (to be published).
 [14] T. V. Kukhto and N. M. Shumeiko, Nucl. Phys. **B219**, 412 (1983); I. V. Akusevich and N. M. Shumeiko, J. Phys. G **20**, 513 (1994).
 [15] Y. S. Tsai, Report No. SLAC-PUB-848, 1971 (unpublished); Y. S. Tsai, Rev. Mod. Phys. **46**, 815 (1974).
 [16] Tables corresponding to Figs. 1 and 2 are available in K. Abe *et al.*, Report No. SLAC-PUB-6734, 1994 (unpublished).
 [17] NMC Collaboration, P. Amaudruz *et al.*, Phys. Lett. B **295**, 159 (1992).
 [18] L. W. Whitlow *et al.*, Phys. Lett. B **250**, 193 (1990).
 [19] L. W. Whitlow *et al.*, Phys. Lett. B **282**, 475 (1992).
 [20] S. J. Brodsky, M. Burkhardt, and I. Schmidt, Report No. SLAC-PUB-6087, 1994 (unpublished).
 [21] R. L. Heimann, Nucl. Phys. **B64**, 429 (1973).
 [22] J. Ellis and M. Karliner, Phys. Lett. B **213**, 73 (1988).
 [23] S. D. Bass and P. V. Landshoff, Phys. Lett. B **336**, 537 (1994); F. E. Close and R. G. Roberts, *ibid.* **336**, 257 (1994).
 [24] M. Lacombe *et al.*, Phys. Rev. C **21**, 861 (1980); M. J. Zuilhof and J. A. Tjon, *ibid.* **22**, 2369 (1980); R. Machleidt *et al.*, Phys. Rep. **149**, 1 (1987).
 [25] Particle Data Group, L. Montanet *et al.*, Phys. Rev. D **50**, 1173 (1994).
 [26] M. S. Schmelling and R. D. St. Denis, Phys. Lett. B **329**, 323 (1994); S. Narison, Report No. CERN-TH-7188/94, 1994 (unpublished).
 [27] F. E. Close and R. G. Roberts, Phys. Rev. Lett. **60**, 1471 (1988).
 [28] J. Ellis and M. Karliner, Phys. Lett. B **341**, 397 (1994), and references therein.